

Line completion number of grid graph $P_n \times P_m$

Joseph Varghese Kureethara^{1*}, Merin Sebastian²

¹Department of Mathematics, Christ University, Bengaluru, India
frjoseph@christuniversity.in

²Department of Mathematics, Christ University, Bengaluru, India
merin.sebastian@maths.christuniversity.in

Received: 19 July 2020; Accepted: 28 January 2021
Published Online: 30 January 2021

Abstract: The concept of super line graph was introduced in the year 1995 by Bagga, Beineke and Varma. Given a graph with at least r edges, the super line graph of index r , $L_r(G)$, has as its vertices the sets of r -edges of G , with two adjacent if there is an edge in one set adjacent to an edge in the other set. The line completion number $lc(G)$ of a graph G is the least positive integer r for which $L_r(G)$ is a complete graph. In this paper, we find the line completion number of grid graph $P_n \times P_m$ for various cases of n and m .

Keywords: Line graph, Super line graph, Grid graph, Line completion number

AMS Subject classification: 05C76

1. Introduction

Line graphs have been studied for over seventy years by now. The line graph transformation is one of the most widely studied of all graph transformations.

In its long history, the concept has been rediscovered several times, with different names such as derived graphs, interchange graphs, edge-to-vertex duals etc. Line graphs can also be considered as intersection graphs. Several variations and generalizations of line graphs have been proposed and studied. These include the concepts of total graphs, path graphs, and others [10].

The *line graph* $L(G)$ of a graph G is defined to have as its vertices the edges of G , with two being adjacent if the corresponding edges share a vertex in G .

* Corresponding Author




Figure 1. Diamond and its line graph

In most generalizations of line graphs, the vertices of the new graph are taken to be another family of subgraphs of G , and adjacency is defined in terms of an appropriate intersection. For example, in the r -path graph $P_r(G)$ of a graph G , the vertices are the paths in G of length r , with adjacency being defined as overlapping in a path of length $r - 1$. We now see an interesting generalization of line graphs.

Given a graph G with at least r edges, the *super line graph (of index r)* $L_r(G)$ has as its vertices the sets of r edges of G , with two adjacent if there is an edge in one set adjacent to an edge in the other set.

Figure 3 is the super line graph of (of index 2) P_5 .




Figure 2. Path P_5




Figure 3. Super Line Graph of index 2, $L_2(P_5)$

2. Line completion number of a graph

Bagga et al. introduced in [8] the concept of super line graph in 1995. Bagga, Beineke and Varma contributed mostly in the study of super line graphs. See Table 1 for a summary the literature.

Authors	Concepts	Ref.	Year
Bagga et al.	super line graphs	[8]	1995
Bagga et al.	properties of super line graphs	[4]	1995
Bagga et al.	line completion number	[7]	1995
Bagga et al.	$L_2(G)$	[6]	1999
Bagga et al.	independence number, pancyclicity	[5]	1999
Gutiérrez and Lladó	edge-residual number, dense graphs	[9]	2002
Bagga et al.	variations and generalizations of line graphs	[1]	2004
Bagga and Ferrero	adjacency matrix, eigenvalues, isolated vertices	[3]	2005
Li et al.	path-comprehensiveness, vertex-pancyclicity	[11]	2008
Bagga et al.	lc of $K_n, P_n, C_n, F_n, W_n, M_n, K_{m,n}$	[2]	2016
Tapadia and Waphare	lc of hypercubes	[12]	2019

Table 1. Summary of the literature on Super Line Graph

Bagga et al. [7] defined the line completion number of a graph as the least index r for which super line graph becomes a complete graph.

One of the important properties of super line graphs is that, if $L_r(G)$ is complete, so is $L_{r+1}(G)$. Bagga et al. determined the line completion numbers for various classes of graphs, viz., complete graph, path, cycle, fan, windmill, wheel, etc. Certain graphs are also characterized with the help of line completion number. In this paper we determine the line completion number of the grid graphs.

3. Line completion number of grid graphs

Let P_n be a path with n vertices and $n - 1$ edges and P_m be a path with m vertices and $m - 1$ edges. The grid graph $P_n \times P_m$ is the Cartesian product of P_n and P_m (See Figure 4). The grid graph has nm vertices and $n(m - 1) + m(n - 1) = 2mn - m - n$ edges. In $P_n \times P_m$, there are n P_m paths and m P_n paths. Let the vertical paths be P_m 's and horizontal paths be P_n 's.

Theorem 1. *Let P_n be a path with n vertices and P_m be a path with m vertices. The*




Figure 4. A general finite grid

line completion number of grid graph $P_n \times P_m$,

$$lc(P_n \times P_m) = \begin{cases} 0 & \text{when } n = m = 1 \\ \lfloor \frac{\max(n,m)}{2} \rfloor & \text{when exactly one of } n \text{ or } m \text{ is } 1 \\ mn + 1 - \frac{(m+n)^2}{2} - \frac{\min(m,n)}{2} & \text{when } n \text{ and } m \text{ are even} \\ mn - \frac{(m+n)^2}{2} - \frac{\min(m,n)+1}{2} & \text{when } n \neq 1 \text{ and } m \neq 1 \text{ are odd} \\ mn + 1 - \min(m,n) - \lceil \frac{\max(m,n)}{2} \rceil & \text{when } n \neq 1 \text{ and } m \neq 1 \\ & \text{are of opposite parity.} \end{cases}$$

Proof. We prove the theorem by analysing all the five cases. Let P_n be the path with n vertices and $n - 1$ edges and P_m be the path with m vertices and $m - 1$ edges. The grid graphs $P_n \times P_m$ and $P_m \times P_n$ are isomorphic graphs.

To find the line completion number of $P_n \times P_m$, our aim is to partition the vertex set of $P_n \times P_m$ into two subsets A and B , so that the subgraphs induced by the two sets A and B , viz., A_k and B_k , respectively, are not adjacent in $P_n \times P_m$. While doing this, we ensure to simultaneously maximize the number of edges in A_k and B_k and minimize the edges that are not in either of these subgraphs. Once we obtain these optimal graphs, adding an edge that is not in A_k or B_k would fetch us the required result.

For the grid graph $P_n \times P_m$, in a Cartesian grid presentation, let the horizontal and the vertical paths be the P_n 's and the P_m 's, respectively.

Case 1. $n = m = 1$.

The graph $P_1 \times P_1$ is P_1 with no edges. Hence, the line completion number trivially is 0.




Figure 5. A path with odd number of vertices



Figure 6. A path with even number of vertices

Case 2. Exactly one of n or m is 1.

The graph in this case is either $P_1 \times P_m$ or $P_n \times P_1$. It is nothing but the path graph. Hence, without loss of generality we consider the graph P_n .

Subcase 2.1. n is odd.

The graph P_n has $n - 1$ edges. Then if we consider a subset of the edge set of P_n with the cardinality ranging from 1 to $\frac{(n-1)}{2} - 1$, there is at least one other subset of the same size that is not adjacent to it. Hence in all such cases, the super line graph is not a complete graph. The two subsets of edges that are not assured to be non-adjacent to each other of the same size are the subsets of consecutive edges from the pendant vertices of the path. See the illustration in Figure 5.

However, if we take a subset of the edge set of size $\frac{(n-1)}{2}$, it is adjacent to every subset of size $\frac{(n-1)}{2}$.

Thus, the line completion number of P_n is $\frac{(n-1)}{2} - 1 + 1 = \frac{(n-1)}{2}$.

Subcase 2.2. n is even.

Continuing in the same line of argument, we can see that the two connected subgraphs each with $\frac{n}{2}$ vertices formed with the two pendant vertices in either of the graphs are non-adjacent. Each of them is of size $\frac{(n-2)}{2}$. See the illustration in Figure 6.

Thus, the line completion number of P_n is $\frac{(n-2)}{2} + 1 = \frac{n}{2}$.

Combining both the cases, we conclude that, the line completion number of a path $P_n \times P_m$ when exactly one of n or m is 1, is $\lfloor \frac{\max(n,m)}{2} \rfloor$. This result is equivalent to the result found in [2].

Slicing of the Grid: Various Scenarios

As indicated in the proof strategy, we need to slice the grid $P_n \times P_m$. In doing so we consider the following cases.

Scenario 1: n is even

Here the vertical slicing of the graph through the central edges of P_n 's will give us two $P_{\frac{n}{2}} \times P_m$'s that are not adjacent to each other and an mK_2 . This is illustrated in the Figure 7.




Figure 7. A vertical slicing




Figure 8. A horizontal slicing

Scenario 2: m is even

If m is even a horizontal slicing of the graph can be done as illustrated in Figure 8. This would generate an nK_2 whose edges are not in the slices.

Scenario 3: n is odd

Since n is odd, there are two central edges for the horizontal P_n 's. Hence, to get two slices of the grid that are equal in size, we do a slicing as presented in Figure 9.

Scenario 4: m is odd

Horizontal slicing of the grid graphs also can be done in an analogous manner. (See Figure 10.)

Scenario 5: m and n are odd

In this case, the number of edges that do not form part of the partition is $m + 3$ and $n + 3$, based on the vertical and horizontal slicing, respectively. See Figure 11 and 12.




Figure 9. An almost vertical slicing




Figure 10. An almost horizontal slicing

Summary of the Analysis

Thus from the five scenarios we analyzed, we come to realize the following.

- R1.** When n is even, the vertical slicing of the grid graph would provide us with two non-adjacent subgraphs, A_p and B_p , of equal size leaving m edges.
- R2.** When m is even, a similar horizontal slicing of the grid graph would provide us with two non-adjacent subgraphs, A_p and B_p , of equal size leaving n edges.
- R3.** When n is odd m is even, an almost vertical slicing of the grid graph would provide us with two non-adjacent subgraphs, A_p and B_p , of equal size leaving $m + 1$ edges.
- R4.** When m is odd and n is even a similar horizontal slicing of the grid graph would




Figure 11. An almost vertical slicing




Figure 12. An almost horizontal slicing

provide us with two non-adjacent subgraphs, A_p and B_p , of equal size leaving $n + 1$ edges.

R5. When n and m are odd, an almost vertical slicing of the grid graph would provide us with two non-adjacent subgraphs, A_p and B_p , of equal size leaving $m + 3$ edges.

R6. When n and m are odd, an almost horizontal slicing of the grid graph would provide us with two non-adjacent subgraphs, A_p and B_p , of equal size leaving $n + 3$ edges.

We now consider non-trivial grids $P_n \times P_m$ with $n > 1$ and $m > 1$.

The grid graph $P_n \times P_m$ has nm vertices and $m(n - 1) + (m - 1)n = 2mn - (m + n)$ edges.

We find the line completion number by considering the various values of n and m .

Case 3. when n and m are even.

We consider two cases.

Subcase 3.1. $m \leq n$.

From the Observation **R1**, we see that a vertical slicing would give use two optimal subgraphs A_p and B_p of size $\frac{2mn-(m+n)-m}{2}$ that are not adjacent to each other and one mK_2 . If we add any edge from mK_2 to any of the subgraphs A_p or B_p , they become adjacent. Hence, the size of A_p = the size B_p = $mn - m - \frac{n}{2}$. If we take the horizontal slicing as given in Observation **R2** leave us with two subgraphs A_p and B_p , each of size $mn - n - \frac{m}{2}$.

Now,

$$mn - n - \frac{m}{2} - (mn - m - \frac{n}{2}) = \frac{m - n}{2}$$

$$\begin{cases} = 0 & \text{if } n = m \\ < 0 & \text{if } n > m. \end{cases}$$

Hence, the vertical slicing gives us more edges. i. e., $mn - m - \frac{n}{2}$ edges. Thus the line completion number of $P_n \times P_m$ in this case is $mn - m - \frac{n}{2} + 1 = mn + 1 - \frac{m+n}{2} - \frac{m}{2}$.

Subcase 3.2. $m \geq n$.

We use the same line of argument as in the previous case and see that the line completion number is obtained from a horizontal slicing. Thus the line completion number of $P_n \times P_m$ is $mn + 1 - \frac{m+n}{2} - \frac{n}{2}$. Since $P_m \times P_n$ is isomorphic to $P_n \times P_m$, we can have a general formula for the line completion number of $P_n \times P_m$. i. e.,

$$lc(P_n \times P_m) = mn + 1 - \frac{m + n}{2} - \frac{\min(m, n)}{2}.$$

Case 4. $n \neq 1$ and $m \neq 1$ are odd.

We consider the following situations.

Subcase 4.1. $m < n$.

From the Observation **R5**, an almost vertical slicing would give use two optimal subgraphs A_p and B_p of equal size leaving exactly $m + 3$ edges. Hence, the size of subgraph A_p is

$$\frac{2mn - (m + n) - (m + 3)}{2} = mn - m - \frac{(n + 3)}{2}.$$

Similarly, an almost horizontal slicing as observed in **R6**, would give us two optimal subgraphs A_p and B_p of equal size leaving exactly $n + 3$ edges. Hence, the size of subgraph A_p is

$$\frac{2mn - (m + n) - (n + 3)}{2} = mn - n - \frac{(m + 3)}{2}.$$

Now,

$$mn - m - \frac{(n+3)}{2} - \left(mn - n - \frac{(m+3)}{2}\right) = \frac{n-m}{2} > 0 \text{ if } n > m.$$

Hence, the vertical slicing gives us more edges. Therefore, the line completion number of the $P_n \times P_m$ in this case is

$$mn - m - \frac{(n+3)}{2} + 1 = mn - \frac{(m+n)}{2} - \frac{(m+1)}{2}.$$

Subcase 4.2. $m > n$.

Since $P_m \times P_n$ is isomorphic to $P_n \times P_m$, we can conclude that,

$$lc(P_n \times P_m) = mn - \frac{(m+n)}{2} - \frac{(n+1)}{2}.$$

A combined formula for this case is thus obtained as,

$$lc(P_n \times P_m) = mn - \frac{(m+n)}{2} - \frac{\min(m,n)+1}{2}.$$

Case 5. when $n \neq 1$ and $m \neq 1$ are of opposite parity.

We consider the following subcases.

Subcase 5.1. $m < n$, m is odd and n is even.

Using Observation **R1**, we do a vertical slicing of the grid graph and obtain two optimal subgraphs each of size

$$mn - m - \frac{n}{2}.$$

Using Observation **R4**, we do an almost horizontal slicing of the grid graph and obtain two subgraphs each of size

$$mn - n - \frac{(m+1)}{2}.$$

However, it is given that $m < n$. Therefore we have,

$$mn - m - \frac{n}{2} - \left(mn - n - \frac{(m+1)}{2}\right) = \frac{(n-m)}{2} + \frac{1}{2} > 0.$$

Thus the vertical slicing provides the larger size. Hence, the line completion number is,

$$\left(mn - m - \frac{n}{2}\right) + 1 = mn + 1 - m - \frac{n}{2}.$$

Subcase 5.2. $m < n$, m is even and n is odd.

Using Observation **R3**, we do an almost vertical slicing of the grid graph and obtain two optimal subgraphs each of size

$$mn - m - \frac{(n + 1)}{2}.$$

Using Observation **R2**, we do a horizontal slicing of the grid graph and obtain two subgraphs each of size

$$mn - n - \frac{m}{2}.$$

However, it is given that $m < n$. Since $n - m$ is at least 1, we have,

$$mn - m - \frac{(n + 1)}{2} - \left(mn - n - \frac{m}{2}\right) = \frac{(n - m)}{2} - \frac{1}{2} \geq 0.$$

Thus the vertical slicing provides the larger size. Hence, the line completion number is,

$$\left(mn - m - \frac{(n + 1)}{2}\right) + 1 = mn + 1 - m - \frac{(n + 1)}{2}.$$

A combined formula for Subcase 5.1 is thus

$$lc(P_n \times P_m) = mn + 1 - m - \lceil \frac{n}{2} \rceil.$$

Subcase 5.3. $m > n$, m is odd and n is even.

Using Observation **R1**, we do a vertical slicing of the grid graph and obtain two optimal subgraphs each of size

$$mn - m - \frac{n}{2}.$$

Using Observation **R4**, we do an almost horizontal slicing of the grid graph and obtain two subgraphs each of size

$$mn - n - \frac{(m + 1)}{2}.$$

As $n - m$ is at most -1 we have,

$$mn - m - \frac{n}{2} - \left(mn - n - \frac{(m + 1)}{2}\right) = \frac{(n - m)}{2} + \frac{1}{2} \leq 0.$$

Thus the vertical slicing provides the larger size. Hence, the line completion number is,

$$\left(mn - n - \frac{(m + 1)}{2}\right) + 1 = mn + 1 - n - \frac{(m + 1)}{2}.$$

Subcase 5.4. $m > n$, m is even and n is odd.

Applying Observation **R3**, we do an almost vertical slicing of the grid graph and obtain two optimal subgraphs, as in Case 3, each of size

$$mn - m - \frac{(n+1)}{2}.$$

Applying Observation **R2**, we do a horizontal slicing of the grid graph and obtain two subgraphs, as in Case 2, each of size

$$mn - n - \frac{(m)}{2}.$$

As $n - m$ is at most -1, we have

$$mn - m - \frac{(n+1)}{2} - \left(mn - n - \frac{m}{2}\right) = \frac{(n-m)}{2} - \frac{1}{2} < 0.$$

Thus the horizontal slicing provides the larger size. Hence, the line completion number is,

$$\left(mn - n - \frac{m}{2}\right) + 1 = mn + 1 - n - \frac{m}{2}.$$

A combined formula for Subcase 5.3 is thus

$$lc(P_n \times P_m) = mn + 1 - n - \lceil \frac{m}{2} \rceil.$$

Subcases 5.1, 5.2, 5.3 and 5.4 can be combined to get the formula for the line completion number of the grid graph when n and m of opposite parity as,

$$lc(P_n \times P_m) = mn + 1 - \min(m, n) - \lceil \frac{\max(m, n)}{2} \rceil.$$

□

4. Illustrations

We see some illustrations now.

Example 1. m and n are even

Consider the graph $P_4 \times P_6$. $P_4 \times P_6$ will have 24 vertices. We partition the vertex set into two equal subsets. Let A and B be the sets. We put vertices of the first 3 P_4 paths in set A . Now the subgraph induced by set A will be $P_4 \times P_3$. Now we take the vertices of 4th, 5th and 6th paths in set B . Set B also induce a subgraph $P_4 \times P_3$. Let C and D be the edge set of $P_4 \times P_3$. From Figure 13, we can see that C and D are not adjacent in $L_{17}(P_4 \times P_6)$. Now, if we add any of the edges $e_{13}, e_{23}, e_{33}, e_{43}$ to set C and D they become adjacent. Therefore the $lc(P_4 \times P_6)$ is 18.




Figure 13. $P_4 \times P_6$

Example 2. $P_4 \times P_5$

Let us consider $P_4 \times P_5$. We partition the vertex set as in Figure 14.




Figure 14. $P_4 \times P_5$

Since both of them are same we prefer the partition in figure 4.3. Now the subgraph induced by these two partitions is $P_2 \times P_5$. Edge set of both copies of $P_2 \times P_5$ will contain 13 edges. Therefore line completion number of $P_4 \times P_5 = 13+1 = 14$.

Example 3. $P_5 \times P_7$

Consider the example $P_5 \times P_7$. Number of vertices in $P_5 \times P_7$ is 35. We cannot partition the vertex set equally. Let A and B be the partitions. Let $|A| = 18$ and $|B| = 17$. Now if we remove the 3rd vertex of 4th P_5 path from set A, we get two equal partitions. Therefore, the number of edges in the subgraphs induced by both A and B = 25. From Figure 15, it is clear that $L_{25}(P_5 \times P_7)$ is not complete. Now if we add one more edge to those corresponding edge sets they become adjacent. Therefore the line completion number is $25+1 = 26$.




Figure 15. $P_5 \times P_7$

Acknowledgement

We express our sincere gratitude to the reviews and comments given by the scholars when we uploaded the manuscript on arxiv.org. Our special thanks to Michel Marcus. We thank the anonymous referees for the comments and corrections provided.

References

- [1] J. Bagga, *Old and new generalizations of line graphs*, Int. J. Math. Math. Sci. **2004** (2004), no. 29, 1509–1521.
- [2] J. Bagga, L. Beineke, and B. Varma, *A number theoretic problem on super line graphs*, AKCE Int. J. Graphs Comb. **13** (2016), no. 2, 177–190.
- [3] J. Bagga, D. Ferrero, and R. Ellis, *The structure of super line graphs*, 8th International Symposium on Parallel Architectures, Algorithms and Networks (IS-PAN'05), IEEE, 2005, pp. 468–471.
- [4] J.S. Bagga, L.W. Beineke, and B.N. Varma, *Super line graphs and their properties*, in: Combinatorics, Graph Theory, Algorithms and Applications, Alavi, Y. and Lick, D. R. and Liu, J. Q., Eds., World Scientific, 1995, pp. 1–5.
- [5] ———, *Independence and cycles in super line graphs*, Australas. J. Combin. **19** (1999), 171–178.
- [6] ———, *The super line graph \mathcal{L}_2* , Discrete Math. **206** (1999), no. 1-3, 51–61.
- [7] K.S. Bagga, L.W. Beineke, and B.N. Varma, *The line completion number of a graph*, in: Graph Theory, Combinatorics, and Applications, Alavi, Y. and Schwen, A., Eds., vol. 2, Wiley-Interscience, New York, 1995, pp. 1197–1201.
- [8] ———, *Super line graphs*, in: Graph Theory, Combinatorics, and Applications, Alavi, Y. and Schwen, A., Eds., vol. 1, Wiley-Interscience, New York, 1995, p. 35–46.

-
- [9] A. Gutiérrez and A.S. Lladó, *On the edge-residual number and the line completion number of a graph*, Ars Combin. **63** (2002), 65–74.
- [10] J. V. Kureethara and J. Kok, *Graphs from subgraphs*, in: Recent Advancements in Graph Theory, Shrimali, N. P. and Shah, N. H., Eds., CRC Press, Boca Raton, 2020, pp. 193–206.
- [11] X. Li, H. Li, and H. Zhang, *Path-comprehensive and vertex-pancyclic properties of super line graph $L_2(G)$* , Discrete Math. **308** (2008), no. 24, 6308–6315.
- [12] S.A. Tapadia and B.N. Waphare, *The line completion number of hypercubes*, AKCE Int. J. Graphs Comb. **16** (2019), no. 1, 78–82.