

On perfectness of annihilating-ideal graph of \mathbb{Z}_n

Manideepa Saha[†], Sucharita Biswas[‡], Angsuman Das^{*}

Department of Mathematics, Presidency University, Kolkata, India

[†]manideepasaha1991@gmail.com

[‡]biswas.sucharita56@gmail.com

^{*}angsuman.maths@presiuniv.ac.in

Received: 25 August 2021; Accepted: 2 December 2021

Published Online: 5 December 2021

Abstract: The annihilating-ideal graph of a commutative ring R with unity is defined as the graph $\mathbb{A}\mathbb{G}(R)$ whose vertex set is the set of all non-zero ideals with non-zero annihilators and two distinct vertices I and J are adjacent if and only if $IJ = 0$. Nikandish *et.al.* proved that $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ is weakly perfect. In this short paper, we characterize n for which $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ is perfect.

Keywords: annihilator, perfect graph, ideals

AMS Subject classification: 05C25, 05C17

1. Introduction

Over the last two decades, various graphs defined on rings have become an interesting topic of research. Various graphs like [1, 2, 4–6, 8, 9] have been constructed to study the interplay between the graph-theoretic and ring-theoretic properties. Interested readers are referred to the following surveys [3, 10, 14] on graphs defined on rings. One such graph is the annihilating-ideal graph $\mathbb{A}\mathbb{G}(R)$ of a commutative ring R , introduced by Behboodi and Rakeei [7].

Definition 1. [7] Let R be a commutative ring with unity. The annihilating-ideal graph of R is defined as the graph $\mathbb{A}\mathbb{G}(R)$ whose vertex set is the set of all non-zero ideals with non-zero annihilators and two distinct vertices I and J are adjacent if and only if $IJ = 0$.

* Corresponding Author

In [15], the authors proved that $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ is *weakly perfect*, i.e., its clique number ω is equal to its chromatic number χ .

A graph G is said to be *perfect* if $\omega(H) = \chi(H)$ holds for all induced subgraphs H of G . Perfect graphs play an important role in graph theory, as many hard graph problems in general like graph coloring, finding maximum clique and independent set, etc. can be solved in polynomial-time in case of perfect graphs. Thus characterizing perfect graphs in different families is an important issue (see [12, 13]). In this short paper, we characterize n for which $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ is perfect. The following theorem is the main result of the paper:

Theorem 1. $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ is perfect if and only if n is one of the forms $p_1^{\alpha_1}$, $p_1^{\alpha_1} p_2^{\alpha_2}$, $p_1^{\alpha_1} p_2 p_3$ or $p_1 p_2 p_3 p_4$, where p_i 's are distinct primes and $\alpha_i \in \mathbb{N}$.

In the next section, we prove Theorem 1. Before that we state an observation and an important result which will be crucial in our proof.

Proposition 1. The vertex set of $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ is $\{\langle m \rangle : m \mid n, 1 < m < n\}$ and two vertices $\langle m_1 \rangle$ and $\langle m_2 \rangle$ are adjacent if and only if $n \mid m_1 m_2$.

Theorem 2. (Strong Perfect Graph Theorem) [11] A graph G is perfect if and only if neither G nor G^c has an induced odd-cycle of length greater or equal to 5.

2. Proof of Theorem 1

We split the proof of Theorem 1 into different cases (lemmas) depending upon the number of distinct prime factors of n .

First, we deal with the case when n has more than 4 distinct prime factors and show that in this case $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ is not perfect.

Lemma 1. If $n = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_k^{\alpha_k}$ and $k \geq 5$, then $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ is not perfect.

Proof. Let $m = n / (p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_5^{\alpha_5})$. Then the following five vertices, taken in order,

$$\langle p_1^{\alpha_1} p_2^{\alpha_2} p_4^{\alpha_4} m \rangle, \langle p_3^{\alpha_3} p_4^{\alpha_4} p_5^{\alpha_5} m \rangle, \langle p_1^{\alpha_1} p_2^{\alpha_2} p_3^{\alpha_3} m \rangle, \langle p_2^{\alpha_2} p_4^{\alpha_4} p_5^{\alpha_5} m \rangle, \langle p_1^{\alpha_1} p_3^{\alpha_3} p_5^{\alpha_5} m \rangle$$

form an induced 5-cycle in $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$. The adjacency and non-adjacency follows from Proposition 1. Hence, by strong perfect graph theorem, the lemma follows. \square

Next we focus on the case when n has exactly 4 distinct prime factors. We characterize the condition when $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ is perfect.

Lemma 2. If $n = p_1^{\alpha_1} p_2^{\alpha_2} p_3^{\alpha_3} p_4^{\alpha_4}$, then $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ is perfect if and only if $\alpha_i = 1$ for all i .

Proof. Let $\alpha_i > 1$ for some i , say $\alpha_1 > 1$. Then the following five vertices, taken in order, $\langle p_1^{\alpha_1} p_4^{\alpha_4} \rangle, \langle p_2^{\alpha_2} p_3^{\alpha_3} p_4^{\alpha_4} \rangle, \langle p_1^{\alpha_1} p_2^{\alpha_2} \rangle, \langle p_1 p_3^{\alpha_3} p_4^{\alpha_4} \rangle, \langle p_1^{\alpha_1-1} p_2^{\alpha_2} p_3^{\alpha_3} \rangle$ form an induced 5-cycle in $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$. As earlier, the adjacency and non-adjacency follows from Proposition 1. Hence, by strong perfect graph theorem, $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ is not perfect. Now, we assume that $n = p_1 p_2 p_3 p_4$. Then $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ has 14 vertices:

1st type:	$\langle p_1 \rangle, \langle p_2 \rangle, \langle p_3 \rangle, \langle p_4 \rangle$	4 vertices of degree 1
2nd type:	$\langle p_1 p_2 \rangle, \langle p_2 p_3 \rangle, \dots, \langle p_3 p_4 \rangle$	6 vertices of degree 3
3rd type:	$\langle p_1 p_2 p_3 \rangle, \langle p_2 p_3 p_4 \rangle, \langle p_1 p_3 p_4 \rangle, \langle p_1 p_2 p_4 \rangle$	4 vertices of degree 7.

If possible, let $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ have an induced odd cycle C of length $t \geq 5$. Thus C must have a vertex of second type. Without loss of generality, let $\langle p_1 p_2 \rangle$ be a vertex in C . As $\langle p_1 p_2 \rangle$ is adjacent to three vertices, namely $\langle p_3 p_4 \rangle, \langle p_1 p_3 p_4 \rangle, \langle p_2 p_3 p_4 \rangle$, at least two of them, must lie on C .

Case 1. $\langle p_1 p_3 p_4 \rangle \sim \langle p_1 p_2 \rangle \sim \langle p_3 p_4 \rangle$ is a part of C .

Let $\langle x \rangle$ be the next vertex on C , i.e., $\langle p_1 p_3 p_4 \rangle \sim \langle p_1 p_2 \rangle \sim \langle p_3 p_4 \rangle \sim \langle x \rangle$. Then by the adjacency condition of the last two vertices, we get $p_1 p_2 \mid x$. But this imply that $\langle x \rangle \sim \langle p_1 p_3 p_4 \rangle$, i.e., we get a chord in C , a contradiction.

Case 2. $\langle p_2 p_3 p_4 \rangle \sim \langle p_1 p_2 \rangle \sim \langle p_3 p_4 \rangle$ is a part of C .

In this case also, proceeding similarly, we get a contradiction.

Case 3. $\langle p_1 p_3 p_4 \rangle \sim \langle p_1 p_2 \rangle \sim \langle p_2 p_3 p_4 \rangle$ is a part of C .

However, in this case, we get a chord of the form $\langle p_1 p_3 p_4 \rangle \sim \langle p_2 p_3 p_4 \rangle$ in C , a contradiction.

Thus $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ has no induced odd cycle C of length $t \geq 5$.

Now, we consider the complement graph of $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$. If possible, let $C' : \langle x_1 \rangle \sim \langle x_2 \rangle \sim \dots \sim \langle x_t \rangle \sim \langle x_1 \rangle$ be an induced odd cycle C' of length $t \geq 5$ in $\mathbb{A}\mathbb{G}^c(\mathbb{Z}_n)$. As C' consists of $t \geq 5$ vertices, at least one of the vertices must be of 1st or 2nd type. We consider the following.

Case 1. $\langle x_1 \rangle$ is a vertex of 1st type, i.e., without loss of generality, let $x_1 = p_1$.

Now, as $\langle p_1 \rangle$ is a pendant vertex in $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$, $\langle p_1 \rangle$ is not adjacent to exactly one vertex in $\mathbb{A}\mathbb{G}^c(\mathbb{Z}_n)$. Thus C' contains a chord, a contradiction.

Case 2. $\langle x_1 \rangle$ is a vertex of 2nd type, i.e., without loss of generality, let $x_1 = p_1 p_2$.

As degree of $\langle p_1 p_2 \rangle$ in $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ is 3, the number of vertices which are not adjacent to $\langle p_1 p_2 \rangle$ in $\mathbb{A}\mathbb{G}^c(\mathbb{Z}_n)$ is 3. Thus, as C' is chordless cycle of length t , x_1 is not adjacent to $(t - 2)$ vertices in C' , i.e., $(t - 2) \leq 3$. Thus C' must be an induced 5-cycle, i.e.,

$$C' : \langle p_1 p_2 \rangle \sim \langle x_2 \rangle \sim \langle x_3 \rangle \sim \langle x_4 \rangle \sim \langle x_5 \rangle \sim \langle p_1 p_2 \rangle.$$

As $\langle x_3 \rangle, \langle x_4 \rangle$ are adjacent to $\langle p_1 p_2 \rangle$ in $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$, we must have $x_3, x_4 \in \{p_3 p_4, p_1 p_3 p_4, p_2 p_3 p_4\}$. If $\{x_3, x_4\} = \{p_1 p_3 p_4, p_2 p_3 p_4\}$, then $\langle x_3 \rangle \sim \langle x_4 \rangle$ in $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$. Thus, without loss of generality, we can assume $x_3 = p_3 p_4$ and $x_4 = p_1 p_3 p_4$, i.e.,

$$C' : \langle p_1 p_2 \rangle \sim \langle x_2 \rangle \sim \langle p_3 p_4 \rangle \sim \langle p_1 p_3 p_4 \rangle \sim \langle x_5 \rangle \sim \langle p_1 p_2 \rangle.$$

As $\langle x_5 \rangle \not\sim \langle p_3 p_4 \rangle$ in $\mathbb{A}\mathbb{G}^c(\mathbb{Z}_n)$, we have $p_1 p_2 \mid x_5$. Thus $x_5 = p_1 p_2 p_3$ or $p_1 p_2 p_4$. However, in any case, $\langle x_5 \rangle \sim \langle p_1 p_3 p_4 \rangle$ in $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$, a contradiction.

Thus $\mathbb{A}\mathbb{G}^c(\mathbb{Z}_n)$ has no induced odd cycle C of length $t \geq 5$. Hence, by strong perfect graph theorem, the lemma follows. \square

Now, we turn towards the case when n has exactly three distinct prime factors and characterize the perfect graphs among this subfamily.

Lemma 3. *If $n = p_1^{\alpha_1} p_2^{\alpha_2} p_3^{\alpha_3}$ and $\alpha_i > 1$ for at least two i 's, then $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ is not perfect.*

Proof. Let $\alpha_i \geq 2$ for at least two i 's, say $\alpha_1, \alpha_2 \geq 2$. Then the following five vertices, taken in order,

$$\langle p_2^{\alpha_2} p_3^{\alpha_3} \rangle, \langle p_1^{\alpha_1} p_2 \rangle, \langle p_1 p_2^{\alpha_2 - 1} p_3^{\alpha_3} \rangle, \langle p_1^{\alpha_1 - 1} p_2^{\alpha_2} \rangle, \langle p_1^{\alpha_1} p_3^{\alpha_3} \rangle$$

form an induced 5-cycle in $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$. As earlier, the adjacency and non-adjacency follows from Proposition 1. Hence, by strong perfect graph theorem, $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ is not perfect. \square

So, now we assume that $n = p_1^{\alpha_1} p_2 p_3$.

Lemma 4. *If $n = p^\alpha q r$, then $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ has no induced odd cycle of length greater than 3.*

Proof. If possible, let $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ has an induced odd cycle $C : \langle x_1 \rangle \sim \langle x_2 \rangle \sim \dots \sim \langle x_t \rangle \sim \langle x_1 \rangle$, where $x_i = p^{\alpha_i} q^{\beta_i} r^{\gamma_i}$ for $i = 1, 2, \dots, t$.

In the next two Claims, we prove that both β_i and γ_i can not be simultaneously 1.

Claim 1. For all $i \in \{1, 2, \dots, t\}$, either $\alpha_i < \alpha/2$ or one of $\beta_i, \gamma_i \neq 1$.

Proof of Claim 1. If possible let $\alpha_i \geq \alpha/2$ and $\beta_i = \gamma_i = 1$. Now $\langle x_i \rangle \sim \langle x_{i+2} \rangle$ and $\langle x_i \rangle \sim \langle x_{i+3} \rangle$ imply $\alpha_{i+2}, \alpha_{i+3} < \alpha/2$, hence $\alpha_{i+2} + \alpha_{i+3} < \alpha$, which is a contradiction as $\langle x_{i+2} \rangle \sim \langle x_{i+3} \rangle$. \blacklozenge

Claim 2. For all $i \in \{1, 2, \dots, t\}$, either $\alpha_i > \alpha/2$ or one of $\beta_i, \gamma_i \neq 1$.

Proof of Claim 2. Without loss of generality let $\alpha_1 \leq \alpha/2$ and $\beta_1 = \gamma_1 = 1$. Now $\langle x_1 \rangle \sim \langle x_2 \rangle$ and $\langle x_1 \rangle \sim \langle x_t \rangle$ imply $\alpha_2, \alpha_t \geq \alpha/2$. As $\langle x_2 \rangle \sim \langle x_t \rangle$ then either $\beta_2 + \beta_t = 0$ or $\gamma_2 + \gamma_t = 0$ or both. Again without loss of generality we can take $\beta_2 + \beta_t = 0$, i.e., $\beta_2 = \beta_t = 0$ and hence $\langle x_2 \rangle \sim \langle x_3 \rangle$ and $\langle x_t \rangle \sim \langle x_{t-1} \rangle$ imply $\beta_3 = 1 = \beta_{t-1}$. Now $\langle x_1 \rangle \sim \langle x_t \rangle$ and $\langle x_1 \rangle \sim \langle x_4 \rangle$ imply $\alpha_1 + \alpha_t \geq \alpha$ and $\alpha_1 + \alpha_4 < \alpha$. From these two equations we have $\alpha_t > \alpha_4$. Therefore $\alpha_3 + \alpha_4 \geq \alpha$ imply $\alpha_3 + \alpha_t > \alpha$. So $\langle x_t \rangle \sim \langle x_3 \rangle$ and $\beta_3 = 1$ imply $\gamma_t + \gamma_3 = 0$, i.e., $\gamma_3 = \gamma_t = 0$. Therefore $\gamma_{t-1} = 1$. As $\langle x_2 \rangle \sim \langle x_{t-1} \rangle$ and $\beta_{t-1} = \gamma_{t-1} = 1$ hence $\alpha_2 + \alpha_{t-1} < \alpha$ and we know $\alpha_1 + \alpha_2 \geq \alpha$. From these two equations we have $\alpha_1 > \alpha_{t-1}$. So $\alpha_{t-1} + \alpha_{t-2} \geq \alpha$ imply $\alpha_1 + \alpha_{t-2} > \alpha$ and $\beta_1 = \gamma_1 = 1$, so we have $\langle x_1 \rangle \sim \langle x_{t-2} \rangle$, which is a contradiction. \blacklozenge

From Claim 1 and Claim 2 we see that for any i , both β_i and γ_i can not be 1. Similarly, it can be shown that both β_i and γ_i can not be 0, because in that case, we must have $\beta_{i+1} = \gamma_{i+1} = 1$, a contradiction.

Claim 3. For all $i \in \{1, 2, \dots, t\}$, $\alpha_i > \alpha/2$.

Proof of Claim 3. Without loss of generality let $\alpha_1 \leq \alpha/2$ and $\beta_1 = 1, \gamma_1 = 0$. Now $\langle x_1 \rangle \sim \langle x_2 \rangle$ and $\langle x_1 \rangle \sim \langle x_t \rangle$ imply $\alpha_2, \alpha_t \geq \alpha/2$ and $\gamma_2 = \gamma_t = 1$. As $\langle x_2 \rangle \approx \langle x_t \rangle$, $\alpha_2 + \alpha_t \geq \alpha$ and $\gamma_2 + \gamma_t = 2$, we have $\beta_2 + \beta_t = 0$, i.e., $\beta_2 = \beta_t = 0$. Hence $\beta_3 = \beta_{t-1} = 1$. Now $\langle x_3 \rangle \approx \langle x_t \rangle$ and $\beta_3 = 1 = \gamma_t$ imply $\alpha_3 + \alpha_t < \alpha$ and $\alpha_2 + \alpha_3 \geq \alpha$, hence $\alpha_2 > \alpha_t$. Therefore $\alpha_t + \alpha_{t-1} \geq \alpha$ implies $\alpha_2 + \alpha_{t-1} > \alpha$. So $\beta_{t-1} = \gamma_2 = 1$ imply $\langle x_2 \rangle \sim \langle x_{t-1} \rangle$, which is impossible and hence the claim holds. \blacklozenge

So from the Claim 1, 2 and 3, we can consider $\alpha_1 > \alpha/2, \beta_1 = 1$ and $\gamma_1 = 0$. Again, from Claim 3 we have $\alpha_3, \alpha_4 > \alpha/2$. So $\langle x_1 \rangle \approx \langle x_3 \rangle$, $\alpha_1 + \alpha_3 > \alpha$ and $\beta_1 = 1$ imply $\gamma_1 + \gamma_3 = 0$, i.e., $\gamma_3 = 0$, i.e., $\gamma_4 = 1$. Therefore $\alpha_1 + \alpha_4 > \alpha$ and $\beta_1 = \gamma_4 = 1$ imply $\langle x_1 \rangle \sim \langle x_4 \rangle$, which is a contradiction. This completes the proof. \square

Lemma 5. If $n = p^\alpha q r$, then $\mathbb{A}\mathbb{G}^c(\mathbb{Z}_n)$ has no induced odd cycle of length greater than 3.

Proof. We start by noting that $\langle a \rangle \sim \langle b \rangle$ in $\mathbb{A}\mathbb{G}^c(\mathbb{Z}_n)$ if and only if $n \nmid ab$. If possible, let $\mathbb{A}\mathbb{G}^c(\mathbb{Z}_n)$ has an induced odd cycle $C : \langle x_1 \rangle \sim \langle x_2 \rangle \sim \dots \sim \langle x_t \rangle \sim \langle x_1 \rangle$, where $x_i = p^{\alpha_i} q^{\beta_i} r^{\gamma_i}$ for $i = 1, 2, \dots, t$.

In the next two Claims, we prove that both β_i and γ_i can not be simultaneously 1.

Claim 1. For all $i \in \{1, 2, \dots, t\}$, either $\alpha_i < \alpha/2$ or one of $\beta_i, \gamma_i \neq 1$.

Proof of Claim 1. If possible let $\alpha_i \geq \alpha/2$ and $\beta_i = \gamma_i = 1$. As $\langle x_{i-1} \rangle$ and $\langle x_{i+1} \rangle \sim \langle x_i \rangle$, hence $\alpha_{i-1}, \alpha_{i+1} < \alpha/2$, i.e., $\alpha_{i-1} + \alpha_{i+1} < \alpha$ and hence $\langle x_{i-1} \rangle \sim \langle x_{i+1} \rangle$, which is a contradiction. \blacklozenge

Claim 2. For all $i \in \{1, 2, \dots, t\}$, either $\alpha_i > \alpha/2$ or one of $\beta_i, \gamma_i \neq 1$.

Proof of Claim 2. Without loss of generality let $\alpha_1 \leq \alpha/2$ and $\beta_1 = \gamma_1 = 1$. As $\langle x_3 \rangle, \langle x_4 \rangle \approx \langle x_1 \rangle$ hence $\alpha_3, \alpha_4 \geq \alpha/2$, i.e., $\alpha_3 + \alpha_4 \geq \alpha$. Now $\langle x_3 \rangle \sim \langle x_4 \rangle$ implies either $\beta_3 + \beta_4 = 0$ or $\gamma_3 + \gamma_4 = 0$ or both. Without loss of generality we can assume $\beta_3 + \beta_4 = 0$, i.e., $\beta_3 = 0 = \beta_4$. Now $\langle x_2 \rangle \approx \langle x_4 \rangle$, $\beta_4 = 0$ imply $\beta_2 = 1$ and $\langle x_3 \rangle \approx \langle x_t \rangle$, $\beta_3 = 0$ imply $\beta_t = 1$. Again $\langle x_1 \rangle \sim \langle x_2 \rangle$ and $\beta_1 = 1 = \gamma_1$ imply $\alpha_1 + \alpha_2 < \alpha$. Therefore $\alpha_2 + \alpha_t \geq \alpha$ implies $\alpha_t > \alpha_1$. So $\alpha_1 + \alpha_{t-1} \geq \alpha$ imply $\alpha_t + \alpha_{t-1} > \alpha$. Now $\langle x_t \rangle \sim \langle x_{t-1} \rangle$ and $\beta_t = 1$ imply $\gamma_t + \gamma_{t-1} = 0$, i.e., $\gamma_t = 0$. Again $\langle x_1 \rangle \sim \langle x_t \rangle$ and $\beta_1 = 1 = \gamma_1$ imply $\alpha_1 + \alpha_t < \alpha$. So $\alpha_2 + \alpha_t \geq \alpha$ imply $\alpha_2 > \alpha_1$. So $\alpha_1 + \alpha_3 \geq \alpha$ imply $\alpha_2 + \alpha_3 > \alpha$. Now $\langle x_2 \rangle \sim \langle x_3 \rangle$ and $\beta_2 = 1$ imply $\gamma_2 + \gamma_3 = 0$, i.e., $\gamma_2 = 0$. Therefore $\gamma_2 = 0 = \gamma_t$ implies $\langle x_2 \rangle \sim \langle x_t \rangle$, which is impossible. \blacklozenge

From Claim 1 and Claim 2 we see that for any i , both β_i and γ_i can not be 1. Similarly, it can be shown that both β_i and γ_i can not be 0, because in that case, we must have $\beta_{i+2} = \gamma_{i+2} = 1$, a contradiction.

Claim 3. For all $i \in \{1, 2, \dots, t\}$, $\alpha_i > \alpha/2$.

Proof of Claim 3. Without loss of generality let $\alpha_1 \leq \alpha/2$ and $\beta_1 = 1, \gamma_1 = 0$. Therefore $\langle x_3 \rangle, \langle x_4 \rangle \approx \langle x_1 \rangle$ imply $\alpha_3, \alpha_4 \geq \alpha/2$ and $\gamma_3 = 1 = \gamma_4$. So $\langle x_3 \rangle \sim \langle x_4 \rangle$

imply $\beta_3 + \beta_4 = 0$, i.e., $\beta_3 = 0 = \beta_4$. Now $\beta_4 = 0$ and $\langle x_4 \rangle \approx \langle x_2 \rangle$ imply $\beta_2 = 1$. Now $\langle x_2 \rangle \sim \langle x_3 \rangle$ and $\beta_2 = 1 = \gamma_3$ imply $\alpha_2 + \alpha_3 < \alpha$. So $\alpha_1 + \alpha_3 \geq \alpha$ imply $\alpha_1 > \alpha_2$. Also $\alpha_2 + \alpha_t \geq \alpha$ imply $\alpha_1 + \alpha_t > \alpha$. So $\langle x_1 \rangle \approx \langle x_t \rangle$ and $\beta_1 = 1$ imply $\gamma_1 + \gamma_t = 0$, i.e., $\gamma_t = 0$, i.e., $\gamma_2 = 1$. Hence we have $\beta_2 = 1 = \gamma_2$, which is not possible by Claim 2. \blacklozenge

So from the Claims 1, 2 and 3, we can consider $\alpha_1 > \alpha/2$, $\beta_1 = 1$ and $\gamma_1 = 0$. So $\langle x_3 \rangle, \langle x_4 \rangle \approx \langle x_1 \rangle$ imply $\gamma_3 = 1 = \gamma_4$. Again from Claim 3 we have $\alpha_3, \alpha_4 > \alpha/2$. So $\langle x_3 \rangle \sim \langle x_4 \rangle$ implies $\beta_3 + \beta_4 = 0$, i.e., $\beta_3 = 0 = \beta_4$. Now $\langle x_2 \rangle \sim \langle x_3 \rangle$, $\gamma_3 = 1$ and $\alpha_2, \alpha_3 > \alpha/2$ imply $\beta_2 + \beta_3 = 0$, i.e., $\beta_2 = 0$. Therefore $\beta_2 = 0 = \beta_4$ imply $\langle x_2 \rangle \sim \langle x_4 \rangle$, which is a contradiction and this completes the proof. \square

Thus, it follows from strong perfect graph theorem and Lemma 4 and Lemma 5, that if $n = p^\alpha q r$, then $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ is perfect.

Thus, the case when n has three distinct prime factors is complete. Now, we focus on the case, when n has two distinct prime factors.

Lemma 6. *If $n = p^\alpha q^\beta$, then $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ has no induced odd cycle of length greater than 3.*

Proof. If possible, let $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ have an induced odd cycle $C : \langle x_1 \rangle \sim \langle x_2 \rangle \sim \dots \sim \langle x_t \rangle \sim \langle x_1 \rangle$, where $x_i = p^{\alpha_i} q^{\beta_i}$ for $i = 1, 2, \dots, t$.

Claim 1. For all $i \in \{1, 2, \dots, t\}$, either $\alpha_i > \alpha/2$ or $\beta_i > \beta/2$.

Proof of Claim 1. If $\alpha_i \leq \alpha/2$ and $\beta_i \leq \beta/2$ for some i , then as $\langle x_i \rangle \sim \langle x_{i+1} \rangle$, we have $\alpha_{i+1} \geq \alpha/2$ and $\beta_{i+1} \geq \beta/2$. Similarly, as $\langle x_i \rangle \sim \langle x_{i-1} \rangle$, we have $\alpha_{i-1} \geq \alpha/2$ and $\beta_{i-1} \geq \beta/2$. But this implies $\alpha_{i+1} + \alpha_{i-1} \geq \alpha$ and $\beta_{i+1} + \beta_{i-1} \geq \beta$, i.e., $\langle x_{i-1} \rangle \sim \langle x_{i+1} \rangle$, a contradiction. Thus the claim holds. \blacklozenge

In Claim 1, we show that for any i , either α_i or β_i is greater than $\alpha/2$ or $\beta/2$ respectively. In the next claim, we show that both of them can not be greater or equal to $\alpha/2$ and $\beta/2$ simultaneously.

Claim 2. For any $i \in \{1, \dots, t\}$, both $\alpha_i \geq \alpha/2$ and $\beta_i \geq \beta/2$ can not hold.

Proof of Claim 2. Without loss of generality, suppose $\alpha_1 \geq \alpha/2$ and $\beta_1 \geq \beta/2$. As $\langle x_1 \rangle \not\sim \langle x_3 \rangle$, we have either $\alpha_1 + \alpha_3 < \alpha$ or $\beta_1 + \beta_3 < \beta$, i.e., $\alpha_3 < \alpha/2$ or $\beta_3 < \beta/2$. Again, without loss of generality, we assume that $\alpha_3 < \alpha/2$. So, by Claim 1, we get $\beta_3 > \beta/2$. As $\langle x_3 \rangle$ is adjacent to both $\langle x_2 \rangle$ and $\langle x_4 \rangle$, we have $\alpha_2, \alpha_4 > \alpha/2$. As $\langle x_1 \rangle \not\sim \langle x_4 \rangle$ and $\alpha_1, \alpha_4 \geq \alpha/2$ and $\beta_1 \geq \beta/2$, we have $\beta_4 < \beta/2$. Again as $\langle x_4 \rangle \sim \langle x_5 \rangle$, we have $\beta_5 > \beta/2$.

Here C is a t -cycle with t odd and $t \geq 5$. We show, by strong induction, that for any odd value of $t \geq 5$, we get a contradiction.

We start with $t = 5$, i.e., $\langle x_1 \rangle \sim \langle x_5 \rangle$. As $\langle x_1 \rangle \not\sim \langle x_4 \rangle$ and $\alpha_1, \alpha_4 \geq \alpha/2$, we have $\beta_1 + \beta_4 < \beta$. Again as $\langle x_4 \rangle \sim \langle x_5 \rangle$, we have $\beta_4 + \beta_5 \geq \beta$. Thus, we get $\beta_5 > \beta_1$. As $\langle x_1 \rangle \sim \langle x_2 \rangle$, we have $\beta_1 + \beta_2 \geq \beta$, i.e., $\beta_2 + \beta_5 > \beta$. Thus as $\langle x_2 \rangle \not\sim \langle x_5 \rangle$, we must have $\alpha_2 + \alpha_5 < \alpha$. Also, as $\langle x_1 \rangle \sim \langle x_5 \rangle$, we have $\alpha_1 + \alpha_5 \geq \alpha$. Thus we must have $\alpha_1 > \alpha_2$. Similarly, $\langle x_2 \rangle \sim \langle x_3 \rangle$ implies $\alpha_2 + \alpha_3 \geq \alpha$, i.e., $\alpha_1 + \alpha_3 > \alpha$. On the other

hand, as $\beta_1, \beta_3 \geq \beta/2$, we have $\beta_1 + \beta_3 \geq \beta$. Thus we have $\langle x_1 \rangle \sim \langle x_3 \rangle$. Hence we get a contradiction for $t = 5$.

For $t > 5$, as $\langle x_1 \rangle \not\sim \langle x_5 \rangle$ and $\beta_1, \beta_5 \geq \beta/2$ and $\alpha_1 \geq \alpha/2$, we have $\alpha_5 < \alpha/2$. Thus the induction hypothesis is: For all odd k satisfying $1 < k < t - 2$, $\alpha_i < \alpha/2$ for odd i with $1 < i \leq k$, $\beta_i \geq \beta/2$ for odd i with $1 \leq i \leq k$, and $\alpha_j \geq \alpha/2$ and $\beta_j < \beta/2$ for even j with $2 < j \leq k - 1$.

Now $\langle x_{k+1} \rangle \sim \langle x_k \rangle$ and $\alpha_k < \alpha/2$ imply $\alpha_{k+1} > \alpha/2$. Similarly, $\langle x_1 \rangle \not\sim \langle x_{k+1} \rangle$ and $\beta_1 \geq \beta/2$ implies $\beta_{k+1} < \beta/2$ and $\langle x_{k+1} \rangle \sim \langle x_{k+2} \rangle$ implies $\beta_{k+2} > \beta/2$. As $k + 2 \leq t - 2$, we have $\langle x_1 \rangle \not\sim \langle x_{k+2} \rangle$ and $\beta_1, \beta_{k+2} > \beta/2$, which implies $\alpha_{k+2} < \alpha/2$. Thus, by induction, we have $\alpha_i < \alpha/2$ for odd i with $1 < i \leq t$, $\beta_i \geq \beta/2$ for odd i , and $\alpha_j \geq \alpha/2$ and $\beta_j < \beta/2$ for even j .

Now, $\langle x_2 \rangle \not\sim \langle x_t \rangle$ implies either $\beta_2 + \beta_t < \beta$ or $\alpha_2 + \alpha_t < \alpha$ or both. As t is odd, $t - 1$ is even and hence $\alpha_1, \alpha_{t-1} \geq \alpha/2$ and $\beta_1 > \beta/2$. Thus $\langle x_1 \rangle \not\sim \langle x_{t-1} \rangle$ implies $\beta_1 + \beta_{t-1} < \beta$ and $\langle x_t \rangle \sim \langle x_{t-1} \rangle$ implies $\beta_t + \beta_{t-1} \geq \beta$. Therefore $\beta_t > \beta_1$.

Again $\langle x_1 \rangle \sim \langle x_2 \rangle$ implies $\beta_1 + \beta_2 \geq \beta$, i.e., $\beta_t + \beta_2 > \beta$. Now, as $\langle x_2 \rangle \not\sim \langle x_t \rangle$, we must have $\alpha_2 + \alpha_t < \alpha$.

Also $\langle x_1 \rangle \sim \langle x_t \rangle$ implies $\alpha_1 + \alpha_t \geq \alpha$. Therefore $\alpha_1 > \alpha_2$. Similarly $\langle x_2 \rangle \sim \langle x_3 \rangle$ implies $\alpha_2 + \alpha_3 \geq \alpha$. Thus $\alpha_1 + \alpha_3 > \alpha$. Again, as $\beta_1, \beta_3 > \beta/2$, we have $\langle x_1 \rangle \sim \langle x_3 \rangle$, a contradiction. Hence Claim 2 holds. \blacklozenge

From Claim 1 and 2, we see that for any i , both α_i, β_i can not be simultaneously ‘greater or equal’ or ‘lesser or equal’ to $\alpha/2$ and $\beta/2$ respectively. So for any i , either $\alpha_i < \alpha/2, \beta_i > \beta/2$ or $\alpha_i > \alpha/2, \beta_i < \beta/2$ holds. Without loss of generality, let $\alpha_1 < \alpha/2$ and $\beta_1 > \beta/2$.

Now, as $\langle x_1 \rangle \sim \langle x_2 \rangle$, we have $\alpha_1 + \alpha_2 \geq \alpha$, which implies $\alpha_2 > \alpha/2$, i.e., $\beta_2 < \beta/2$ (by Claim 2). Similarly $\langle x_2 \rangle \sim \langle x_3 \rangle$ implies $\beta_3 > \beta/2$, i.e., $\alpha_3 < \alpha/2$ (by Claim 2). Proceeding this way, we get $\alpha_i < \alpha/2$ and $\beta_i > \beta/2$ when i is odd and, $\alpha_i > \alpha/2$ and $\beta_i < \beta/2$ if i is even.

As t is odd, we have $\alpha_t < \alpha/2$. Also, as $\langle x_1 \rangle \sim \langle x_t \rangle$, we have $\alpha_1 + \alpha_t \geq \alpha$. However as $\alpha_1, \alpha_t < \alpha/2$, we get a contradiction. Thus $\mathbb{A}\mathbb{G}(\mathbb{Z}_n)$ has no induced odd cycle of length greater than 3. \square

Lemma 7. *If $n = p^\alpha q^\beta$, then $\mathbb{A}\mathbb{G}^c(\mathbb{Z}_n)$ has no induced odd cycle of length greater than 3.*

Proof. We start by noting that $\langle a \rangle \sim \langle b \rangle$ in $\mathbb{A}\mathbb{G}^c(\mathbb{Z}_n)$ if and only if $n \nmid ab$. If possible, let $\mathbb{A}\mathbb{G}^c(\mathbb{Z}_n)$ has an induced odd cycle $C : \langle x_1 \rangle \sim \langle x_2 \rangle \sim \dots \sim \langle x_t \rangle \sim \langle x_1 \rangle$, where $x_i = p^{\alpha_i} q^{\beta_i}$ for $i = 1, 2, \dots, t$.

Claim 1. For all $i \in \{1, 2, \dots, t\}$, either $\alpha_i > \alpha/2$ or $\beta_i > \beta/2$.

Proof of Claim 1. If $\alpha_i \leq \alpha/2$ and $\beta_i \leq \beta/2$ for some i , then as $\langle x_i \rangle \not\sim \langle x_{i+2} \rangle$ and $\langle x_i \rangle \not\sim \langle x_{i+3} \rangle$, we have $\alpha_{i+2}, \alpha_{i+3} \geq \alpha/2$ and $\beta_{i+2}, \beta_{i+3} \geq \beta/2$. But this imply that $\langle x_{i+2} \rangle \not\sim \langle x_{i+3} \rangle$ in $\mathbb{A}\mathbb{G}^c(\mathbb{Z}_n)$, a contradiction. Hence Claim 1 holds. \blacklozenge

In Claim 1, we show that for any i , either α_i or β_i is greater than $\alpha/2$ or $\beta/2$ respectively. In the next claim, we show that both of them can not be greater or equal to $\alpha/2$ and $\beta/2$ simultaneously.

Claim 2. For any i , both $\alpha_i \geq \alpha/2$ and $\beta_i \geq \beta/2$ can not hold.

Proof of Claim 2. Without loss of generality, suppose $\alpha_1 \geq \alpha/2$ and $\beta_1 \geq \beta/2$. $\langle x_1 \rangle \sim \langle x_2 \rangle$ implies either $\alpha_1 + \alpha_2 < \alpha$ or $\beta_1 + \beta_2 < \beta$ or both. Again, without loss of generality, we assume that $\alpha_1 + \alpha_2 < \alpha$, i.e., $\alpha_2 < \alpha/2$. Now $\langle x_2 \rangle \approx \langle x_t \rangle$ implies $\alpha_2 + \alpha_t \geq \alpha$, i.e., $\alpha_t > \alpha/2$.

At first we assume that $t = 5$. Therefore $\langle x_1 \rangle \sim \langle x_5 \rangle$ and $\alpha_1, \alpha_5 \geq \alpha/2$ imply $\beta_1 + \beta_5 < \beta$. Now $\langle x_1 \rangle \approx \langle x_3 \rangle$ imply $\alpha_1 + \alpha_3 \geq \alpha$, so $\alpha_1 + \alpha_2 < \alpha$ implies $\alpha_3 > \alpha_2$. Again $\langle x_3 \rangle \sim \langle x_4 \rangle$ imply either $\alpha_3 + \alpha_4 < \alpha$ or $\beta_3 + \beta_4 < \beta$ or both. Now $\langle x_2 \rangle \approx \langle x_4 \rangle$ imply $\alpha_2 + \alpha_4 \geq \alpha$. If $\alpha_3 + \alpha_4 < \alpha$, then we have $\alpha_2 > \alpha_3$, which is a contradiction as we already have $\alpha_3 > \alpha_2$. Now $\langle x_1 \rangle \approx \langle x_4 \rangle$ implies $\beta_1 + \beta_4 \geq \beta$. If $\beta_3 + \beta_4 < \beta$, then we have $\beta_1 > \beta_3$. Now $\langle x_3 \rangle \approx \langle x_5 \rangle$ implies $\beta_3 + \beta_5 \geq \beta$, therefore $\beta_1 + \beta_5 > \beta$, which contradicts the condition $\beta_1 + \beta_5 < \beta$. So for $t = 5$ the Claim 2 is true.

Now assume that $t > 5$. As $\alpha_1, \alpha_t \geq \alpha/2$ and $\langle x_1 \rangle \sim \langle x_t \rangle$, we have $\beta_1 + \beta_t < \beta$, i.e., $\beta_t < \beta/2$ as $\beta_1 \geq \beta/2$. Now $\alpha_2 < \alpha/2$ and $\langle x_4 \rangle, \langle x_5 \rangle \approx \langle x_2 \rangle$ imply $\alpha_4, \alpha_5 > \alpha/2$, hence $\alpha_4 + \alpha_5 > \alpha$. Again $\beta_t < \beta/2$ and $\langle x_4 \rangle, \langle x_5 \rangle \approx \langle x_t \rangle$ imply $\beta_4, \beta_5 > \beta/2$, hence $\beta_4 + \beta_5 > \beta$, which is a contradiction as $\langle x_4 \rangle \sim \langle x_5 \rangle$. Hence Claim 2 holds for all odd $t \geq 5$. \blacklozenge

From Claims 1 and 2, we see that for any i , both α_i, β_i can not be simultaneously ‘greater or equal’ or ‘lesser or equal’ to $\alpha/2$ and $\beta/2$, respectively. So for any i , either $\alpha_i < \alpha/2, \beta_i > \beta/2$ or $\alpha_i > \alpha/2, \beta_i < \beta/2$ holds. Without loss of generality, let $\alpha_1 < \alpha/2$ and $\beta_1 > \beta/2$.

Now $\langle x_3 \rangle, \langle x_4 \rangle \approx \langle x_1 \rangle$ imply $\alpha_3, \alpha_4 > \alpha/2$ and hence by Claim 2 we have $\beta_3, \beta_4 < \beta/2$. As $\langle x_2 \rangle \approx \langle x_4 \rangle$, so $\beta_2 > \beta/2$ and by Claim 2 we have $\alpha_2 < \alpha/2$. Now $\langle x_2 \rangle \approx \langle x_5 \rangle$ implies $\alpha_5 > \alpha/2$. Then by Claim 2 we have $\beta_5 < \beta/2$, but $\beta_3 < \beta/2$ imply $\beta_3 + \beta_5 < \beta$, which is a contradiction as $\langle x_3 \rangle \approx \langle x_5 \rangle$. Thus $\text{AG}^c(\mathbb{Z}_n)$ has no induced odd cycle of length greater than 3. \square

Thus from Lemma 6 and Lemma 7, we have if $n = p^\alpha q^\beta$, then $\text{AG}(\mathbb{Z}_n)$ is perfect.

Now, we deal with the last case when n is a prime power.

Lemma 8. *If $n = p^\alpha$, then $\text{AG}(\mathbb{Z}_n)$ is perfect.*

Proof. In this case, the vertices are $\langle p \rangle, \langle p^2 \rangle, \dots, \langle p^{\alpha-1} \rangle$ and two vertices $\langle p^k \rangle$ and $\langle p^l \rangle$ are adjacent in $\text{AG}(\mathbb{Z}_n)$ if and only if $k + l \geq \alpha$.

If possible, let $C : \langle p^{k_1} \rangle \sim \langle p^{k_2} \rangle \sim \dots \sim \langle p^{k_t} \rangle \sim \langle p^{k_1} \rangle$ be an induced odd cycle of length $t \geq 5$. By adjacency condition, we have $k_i + k_{i+1} \geq \alpha$ for each $i \in \{1, \dots, t\}$, where $k_{t+1} = k_1$, and hence $2(k_1 + k_2 + \dots + k_t) \geq t\alpha$. On the other hand, from non-adjacency condition we have $k_i + k_{i+2} < \alpha$ for each $i \in \{1, \dots, t\}$, where $k_{t+2} = k_2$, and so $2(k_1 + k_2 + \dots + k_t) \geq t\alpha$ which is a contradiction. Thus $\text{AG}(\mathbb{Z}_n)$ has no induced odd cycle C of length $t \geq 5$. Proceeding similarly, it can be shown that $\text{AG}^c(\mathbb{Z}_n)$ also has no induced odd cycle of length $t \geq 5$. Hence $\text{AG}(\mathbb{Z}_n)$ is perfect. \square

Combining all the results in this section, we get the proof of Theorem 1.

Acknowledgement

The first and third authors acknowledge the funding of DST-SERB-SRG Sanction no. *SRG/2019/000475* and *SRG/2019/000684*, Govt. of India. The second author is supported by the PhD fellowship of CSIR (File no. 08/155(0086)/2020-EMR-I), Govt. of India.

References

- [1] C. AbdIoglu, E.Y. Cellkel, and A. Das, *The armendariz graph of a ring*, Discuss. Math. Gen. Algebra Appl. **38** (2018), no. 2, 189–196.
- [2] J. Amjadi, R. Khoeilar, and A. Alilou, *The annihilator-inclusion ideal graph of a commutative ring*, Commun. Comb. Optim. **6** (2021), no. 2, 231–248.
- [3] D.F. Anderson, M.C. Axtell, and J.A. Stickles, Jr., *Zero-divisor graphs in commutative rings*, Commutative Algebra: Noetherian and Non-Noetherian Perspectives (M. Fontana, S.-E. Kabbaj, B. Olberding, and I. Swanson, eds.), Springer, 2011, pp. 23–45.
- [4] D.F. Anderson and A. Badawi, *The total graph of a commutative ring*, J. Algebra **320** (2008), no. 7, 2706–2719.
- [5] D.F. Anderson and P.S. Livingston, *The zero-divisor graph of a commutative ring*, J. Algebra **217** (1999), no. 2, 434–447.
- [6] A. Badawi, *On the annihilator graph of a commutative ring*, Comm. Algebra **42** (2014), no. 1, 108–121.
- [7] M. Behboodi and Z. Rakeei, *The annihilating-ideal graph of commutative rings I*, J. Algebra . Appl. **10** (2011), no. 4, 727–739.
- [8] B. Bose and A. Das, *Graph theoretic representation of rings of continuous functions*, Filomat **34** (2020), no. 10, 3417–3428.
- [9] I. Chakrabarty, S. Ghosh, T.K. Mukherjee, and M.K. Sen, *Intersection graphs of ideals of rings*, Discrete Math. **309** (2009), no. 17, 5381–5392.
- [10] I. Chakrabarty and J.V. Kureethara, *A survey on the intersection graphs of ideals of rings*, Commun. Comb. Optim. (in press).
- [11] M. Chudnovsky, N. Robertson, P. Seymour, and R. Thomas, *The strong perfect graph theorem*, Annals of Math. **164** (2006), no. 1, 51–229.
- [12] A. Das, *On perfectness of intersection graph of ideals of \mathbb{Z}_n* , Discuss. Math. Gen. Algebra Appl. **37** (2017), no. 2, 119–126.
- [13] M. Ebrahimi, *The character graph of a finite group is perfect*, Bull. Aust. Math. Soc. **104** (2021), no. 1, 127–131.
- [14] H.R. Maimani, M.R. Pournaki, A. Tehranian, and S. Yassemi, *Graphs attached to rings revisited*, Arab. J. Sci. Eng. **36** (2011), no. 6, 997–1011.
- [15] R. Nikandish, H.R. Maimani, and H. Izanloo, *The annihilating-ideal graph of \mathbb{Z}_n is weakly perfect*, Contrib. Discrete Math. **11** (2016), no. 1, 16–21.