

## Tetravalent half-arc-transitive graphs of order $12p$

M. Ghasemi<sup>1,\*</sup>, N. Mehdipoor<sup>2,†</sup> and A.A. Talebi<sup>3,‡</sup>

<sup>1</sup>Department of Mathematics, Urmia University, Urmia 57135, Iran  
\*m.ghasemi@urmia.ac.ir

<sup>2</sup>Department of Mathematics, University of Mazandaran, Babolsar, Iran  
†nargesmehdipoor@yahoo.com  
‡a.talebi@umz.ac.ir

*Received: 8 September 2022; Accepted: 16 November 2022*  
*Published Online: 20 November 2022*

**Abstract:** A graph is half-arc-transitive if its automorphism group acts transitively on its vertex set, edge set, but not its arc set. In this paper, we study all tetravalent half-arc-transitive graphs of order  $12p$ , where  $p$  is a prime.

**Keywords:** Half-arc-transitive graph, Tightly attached, Regular covering projection, Solvable groups

**AMS Subject classification:** 05C25, 20B25

### 1. Introduction

In this study, all graphs considered are assumed to be finite, simple and connected. For a graph  $X$ ,  $V(X)$ ,  $E(X)$ ,  $A(X)$  and  $\text{Aut}(X)$  denote its vertex set, edge set, arc set, and full automorphism group, respectively. For  $u, v \in V(X)$ ,  $\{u, v\}$  denotes the edge incident to  $u$  and  $v$  in  $X$ , and  $N_X(u)$  denotes the neighborhood of  $u$  in  $X$ , that is, the set of vertices adjacent to  $u$  in  $X$ .

A graph  $\tilde{X}$  is called a covering of a graph  $X$  with projection  $p: \tilde{X} \rightarrow X$  if there is a surjection  $p: V(\tilde{X}) \rightarrow V(X)$  such that  $p|_{N_{\tilde{X}}(\tilde{v})}: N_{\tilde{X}}(\tilde{v}) \rightarrow N_X(v)$  is a bijection for any vertex  $v \in V(X)$  and  $\tilde{v} \in p^{-1}(v)$ . A permutation group  $G$  on a set  $\Omega$  is said to be semiregular if the stabilizer  $G_v$  of  $v$  in  $G$  is trivial for each  $v \in \Omega$ , and is regular if  $G$  is transitive, and semiregular. Let  $K$  be a subgroup of  $\text{Aut}(X)$  such that  $K$  is intransitive on  $V(X)$ . The quotient graph  $X/K$  induced by  $K$  is defined as the graph

---

\* Corresponding Author

such that the set  $\Omega$  of  $K$ -orbits in  $V(X)$  is the vertex set of  $X/K$  and  $B, C \in \Omega$  are adjacent if and only if there exists a  $u \in B$  and  $v \in C$  such that  $\{u, v\} \in E(X)$ . A covering  $\tilde{X}$  of  $X$  with a projection  $p$  is said to be regular (or  $K$ -covering) if there is a subgroup  $K$  of the automorphism group  $\text{Aut}(\tilde{X})$  such that  $K$  is semiregular on both  $V(\tilde{X})$  and  $E(\tilde{X})$  and graph  $X$  is isomorphic to the quotient graph  $\tilde{X}/K$ , say by  $h$ , and the quotient map  $\tilde{X} \rightarrow \tilde{X}/K$  is the composition  $ph$  of  $p$  and  $h$ . The group of covering transformations  $\text{CT}(p)$  of  $p: \tilde{X} \rightarrow X$  is the group of all self equivalences of  $p$ , that is, of all automorphisms  $\tilde{\alpha} \in \text{Aut}(\tilde{X})$  such that  $p = \tilde{\alpha}p$ . If  $\tilde{X}$  is connected,  $K$  becomes the covering transformation group.

For a graph  $X$  and a subgroup  $G$  of  $\text{Aut}(X)$ ,  $X$  is said to be  $G$ -vertex-transitive,  $G$ -edge-transitive or  $G$ -arc-transitive if  $G$  is transitive on  $V(X)$ ,  $E(X)$  or  $A(X)$ , respectively, and  $G$ -arc-regular if  $G$  acts regularly on  $A(X)$ . A graph  $X$  is called vertex-transitive, edge-transitive, arc-transitive, or arc-regular if  $X$  is  $\text{Aut}(X)$ -vertex-transitive,  $\text{Aut}(X)$ -edge-transitive,  $\text{Aut}(X)$ -arc-transitive, or  $\text{Aut}(X)$ -arc-regular, respectively. Let  $X$  be a tetravalent  $G$ -half-arc-transitive graph for a subgroup  $G$  of  $\text{Aut}(X)$ , that is  $G$  acts transitively on  $V(X)$ ,  $E(X)$ , but not  $A(X)$ . Then under the natural action of  $G$  on  $V(X) \times V(X)$ ,  $G$  has two orbits on the arc set  $A(X)$ , say  $A_1$  and  $A_2$ , where  $A_2 = \{(v, u) | (u, v) \in A_1\}$ . Therefore, one may obtain two oriented graphs with the vertex set  $V(X)$  and the arc sets  $A_1$  and  $A_2$ . Assume that  $D_G(X)$  be one of the two oriented graphs. Also in the special case, if  $G = \text{Aut}(X)$  then  $X$  is said to be  $1/2$ -transitive or half-arc-transitive.

By Tutte [29], each connected vertex-transitive and edge-transitive graph of odd valency is arc-transitive. So half-arc-transitive graphs of odd valency do not exist. Bouwer [5] answered Tutte's question about existence of half-arc-transitive graphs of even valency. A number of authors later studied the construction of these graphs. See, for example [1, 2, 9, 11, 14, 20–23, 30, 33, 36]. Let  $p$  be a prime. There are no half-arc-transitive graphs of order  $p$ ,  $p^2$  and  $2p$  (see [6, 8]). Feng, Kwak, Wang and Zhou [12] classified the connected tetravalent half-arc-transitive graphs of order  $2pq$  for distinct odd primes  $p$  and  $q$ . The tetravalent half-arc-transitive graphs of order  $p^5$ ,  $p^4$ ,  $2p^2$ ,  $p^3$  and  $2p^3$  are classified in [7, 13, 34, 37, 38] respectively. Wang et al. [32] studied tetravalent half-arc-transitive graphs of order a product of three primes. In [24], Liu studied tetravalent half-arc-transitive graphs of order  $p^2q^2$  with  $p, q$  distinct odd primes. Feng et al. [15] classified the tetravalent half-arc-transitive graphs of order  $4p$ . In [10] a complete classification of tetravalent half-arc-transitive metacirculants of order 2-powers was given. In [35], a classification of all tetravalent half-arc-transitive graphs of order  $8p$  was given. In this paper, we will study tetravalent half-arc-transitive graphs of order  $12p$ .

## 2. Preliminaries

Let  $X$  be a graph and  $K$  be a finite group. By  $a^{-1}$  we mean the reverse arc to an arc  $a$ . A voltage assignment (or  $K$ -voltage assignment) of  $X$  is a function  $\xi: A(X) \rightarrow K$  with the property that  $\xi(a^{-1}) = \xi(a)^{-1}$  for each arc  $a \in A(X)$ . The values of  $\xi$  are

called voltages, and  $K$  is the voltage group. The graph  $X \times_{\xi} K$  derived from a voltage assignment  $\xi : A(X) \rightarrow K$  has vertex set  $V(X) \times K$  and edge set  $E(X) \times K$ , so that an edge  $(e, g)$  of  $X \times K$  joins a vertex  $(u, g)$  to  $(v, \xi(a)g)$  for  $a = (u, v) \in A(X)$  and  $g \in K$ , where  $e = \{u, v\}$ . Clearly, the derived graph  $X \times_{\xi} K$  is a covering of  $X$  with the first coordinate projection  $p : X \times_{\xi} K \rightarrow X$ , which is called the natural projection. By defining  $(u, g')^g = (u, g'g)$  for any  $g \in K$  and  $(u, g') \in V(X \times_{\xi} K)$ ,  $K$  becomes a subgroup of  $\text{Aut}(X \times_{\xi} K)$  which acts semiregularly on  $V(X \times_{\xi} K)$ . Therefore,  $X \times_{\xi} K$  can be viewed as a  $K$ -covering. For each  $u \in V(X)$  and  $\{u, v\} \in E(X)$ , the vertex set  $\{(u, g) | g \in K\}$  is the fibre of  $u$  and the edge set  $\{(u, g)(v, \xi(a)g) | g \in K\}$  is the fibre of  $\{u, v\}$ , where  $a = (u, v)$ . The group  $K$  of automorphisms of  $X$  fixing every fibre setwise is called the covering transformation group. Conversely, each regular covering  $\tilde{X}$  of  $X$  with a covering transformation group  $K$  can be derived from a  $K$ -voltage assignment. Given a spanning tree  $T$  of the graph  $X$ , a voltage assignment  $\xi$  is said to be  $T$ -reduced if the voltages on the tree arcs are the identity. Gross and Tucker in [18] showed that every regular covering  $\tilde{X}$  of a graph  $X$  can be derived from a  $T$ -reduced voltage assignment  $\tilde{\xi}$  with respect to an arbitrary fixed spanning tree  $T$  of  $X$ .

Let  $\tilde{X}$  be a  $K$ -covering of  $X$  with a projection  $p$ . If  $\alpha \in \text{Aut}(X)$  and  $\tilde{\alpha} \in \text{Aut}(\tilde{X})$  satisfy  $\tilde{\alpha}p = p\alpha$ , we call  $\tilde{\alpha}$  a lift of  $\alpha$ , and  $\alpha$  the projection of  $\tilde{\alpha}$ . The lifts and projections of such subgroups are of course subgroups in  $\text{Aut}(\tilde{X})$  and  $\text{Aut}(X)$ , respectively.

Let  $G$  be a group, and let  $S \subseteq G$  be a set of group elements such that the identity element 1 not in  $S$ . The Cayley graph associated with  $(G, S)$  is defined as the graph having one vertex associated with each group element, edges  $(g, h)$  whenever  $hg^{-1}$  in  $S$ . The Cayley graph  $X$  is denoted by  $\text{Cay}(G, S)$ . In graph theory, the lexicographic product or (graph) composition  $G[H]$  of graphs  $G$  and  $H$  is a graph such that the vertex set of  $G[H]$  is the cartesian product  $V(G) \times V(H)$ ; and any two vertices  $(x, y)$  and  $(v, w)$  are adjacent in  $G[H]$  if and only if either  $x$  is adjacent with  $v$  in  $G$  or  $v = x$  and  $w$  is adjacent with  $y$  in  $H$ . Clearly, if  $G$  and  $H$  are arc-transitive then  $G[H]$  is arc-transitive.

Let  $X$  be a tetravalent  $G$ -half-arc-transitive graph for some  $G \leq \text{Aut}(X)$ . Then no element of  $G$  can interchange a pair of adjacent vertices in  $X$ . By [19], there is no half-arc-transitive graph with less than 27 vertices. Half-arc-transitive graphs have even valencies. An even length cycle  $C$  in  $X$  is a  $G$ -alternating cycle if every other vertex of  $C$  is the head and every other vertex of  $C$  is the tail of its two incident edges in  $D_G(X)$ . All  $G$ -alternating cycles in  $X$  have the same length. The radius of graph is half of the length of an alternating cycle. Any two adjacent  $G$ -alternating cycles in  $X$  intersect in the same number of vertices, called the  $G$ -attachment number of  $X$ . The intersection of two adjacent  $G$ -alternating cycles is called a  $G$ -attachment set. We say that  $X$  is tightly attached if the attachment number of  $X$  equal with its radius.

Now we introduce graph  $X(r; m, n)$  and a result due to Marušič.

Suppose that  $m \geq 3$  be an integer,  $n \geq 3$  an odd integer and let  $r \in \mathbb{Z}_n^*$  satisfy  $r^m = \pm 1$ . The graph  $X(r; m, n)$  is defined to have vertex set  $V = \{u_i^j \mid i \in \mathbb{Z}_m, j \in$

$\mathbb{Z}_n\}$  and edge set  $E = \{\{u_i^j, u_{i+1}^{j\pm r^i}\} \mid i \in \mathbb{Z}_m, j \in \mathbb{Z}_n\}$ .

**Proposition 1.** [25, Theorem 3.4] *A connected tetravalent graph  $X$  is a tightly attached half-arc-transitive graph of odd radius  $n$  if and only if  $X \cong X(r; m, n)$ , where  $m \geq 3$ , and  $r \in \mathbb{Z}_n^*$  satisfying  $r^m = \pm 1$ , and moreover none of the following conditions is fulfilled:*

- (1)  $r^2 = \pm 1$ ;
- (2)  $(r; m, n) = (2; 3, 7)$ ;
- (3)  $(r; m, n) = (r; 6, 7k)$ , where  $k \geq 1$  is odd,  $(7, k) = 1$ ,  $r^6 = 1$ , and there exists a unique solution  $q \in \{r, -r, r^{-1}, -r^{-1}\}$  of the equation  $x^2 + x - 2 = 0$  such that  $7(q - 1) = 0$  and  $q \equiv 5 \pmod{7}$ .

The following is the main result of the paper tetravalent half-transitive graphs of order  $4p$ .

**Proposition 2.** [15, Theorem 3.3] *Let  $p$  be a prime and  $X$  a tetravalent graph of order  $4p$ . Then,  $X$  is half-transitive if and only if  $p \equiv 1 \pmod{8}$  and  $X \cong X(r; 4, p)$  (denote by  $X(4, p)$  the graph  $X(r; 4, p)$ ).*

Now we express an observations about tetravalent half-arc-transitive graphs.

**Proposition 3.** [26, Lemma 3.5] *Let  $X$  be a connected tetravalent  $G$ -half-arc-transitive graph for some  $G \leq \text{Aut}(X)$ , and let  $\Delta$  be a  $G$ -attachment set of  $X$ . If  $|\Delta| \geq 3$ , then the vertex-stabilizer of  $v \in V(X)$  in  $G$  is of order 2.*

**Proposition 4.** [17] *A non-abelian simple group whose order has at most three prime divisors is isomorphic to one of the following groups:*

$$A_5, A_6, \text{PSL}(2, 7), \text{PSL}(2, 8), \text{PSL}(2, 17), \text{PSL}(3, 3), \text{PSU}(3, 3), \text{PSU}(4, 2),$$

*whose orders are  $2^2 \times 3 \times 5$ ,  $2^3 \times 3^2 \times 5$ ,  $2^3 \times 3 \times 7$ ,  $2^3 \times 3^2 \times 7$ ,  $2^4 \times 3^2 \times 17$ ,  $2^4 \times 3^3 \times 13$ ,  $2^5 \times 3^3 \times 7$ ,  $2^6 \times 3^4 \times 5$ , respectively.*

The following result is extracted from [4, Theorem 1].


**Proposition 5.** *Let  $X$  be a tetravalent arc-transitive graph of order  $2pq$  where  $p$  and  $q$  are odd and distinct primes. Then one of the following holds:*

- (1)  $X$  is arc-regular and appears in [40];
- (2)  $X$  is isomorphic to the lexicographic product  $C_{pq}[2K_1]$  of the cycle  $C_{pq}$  and the edgeless graph on two vertices  $2K_1$ .


In the following, we describe the structure of the graphs required in this paper [[27], [28], [39]].

The Rose Window graph  $R_6(5, 4)$  is a tetravalent graph with 12 vertices. Its vertex set is  $\{S_i, Q_i \mid i \in \mathbb{Z}_6\}$ . The graph has four kinds of edges: kind of edges:  $S_i S_{i+1}$  (rim edges),  $S_i Q_i$  (inspoke edges),  $S_{i+5} Q_i$  (outspoke edges) and  $Q_i Q_{i+4}$  (hub edges).  $|\text{Aut}(R_6(5, 4))| = 48$ . Figure 1 shows  $R_6(5, 4)$ .

A general Wreath graph  $W(6, 2)$  has 12 vertices and it is regular of valency 4. Its vertex set is  $\{E_i, F_i | i \in \mathbb{Z}_6\}$ , where  $E_i = (i, 0)$  and  $F_i = (i, 1)$ . Its edges are  $\{E_i, E_{i+1}\}$ ,  $\{E_i, F_{i+1}\}$ ,  $\{F_i, E_{i+1}\}$  and  $\{F_i, F_{i+1}\}$ .  $|\text{Aut}(W(6, 2))| = 768$ . See Figure 2.



**Figure 1.** The Rose Window graph  $R_6(5, 4)$



**Figure 2.** The Wreath graph  $W(6, 2)$

The graph  $C(2; p, 2)$  was first defined by Praeger and Xu [28, Definition 2.1 (b)]. Let  $p$  be an odd prime. The graph  $C(2; p, 2)$  has vertex set  $\mathbb{Z}_p \times (\mathbb{Z}_2 \times \mathbb{Z}_2)$  and its edges are defined by  $\{(i, (x, y)), (i + 1, (y, z))\} \in E(C(2; p, 2))$  for all  $i \in \mathbb{Z}_p$  and  $x, y, z \in \mathbb{Z}_2$ .  $\text{Aut}(C(2; p, 2)) \cong D_{2p} \times \mathbb{Z}_2^p$ .

Let  $p \equiv 1 \pmod{4}$ , where  $p$  is a prime and  $w$  is an element of order 4 in  $\mathbb{Z}_p^*$ . The graph  $CA_{4p}^0$  is  $\text{Cay}(G, \{a, a^{-1}, a^{w^2}b, a^{-w^2}b\})$  and the graph  $CA_{4p}^1$  is  $\text{Cay}(G, \{a, a^{-1}, a^wb, a^{-wb}\})$ , where  $G = \langle a \rangle \times \langle b \rangle \cong \mathbb{Z}_{2p} \times \mathbb{Z}_2$ .

### 3. Main Results

In this section, we study all tetravalent half-arc-transitive graphs of order  $12p$  where  $p$  is a prime. To do this, we prove the following results.

**Lemma 1.** *Let  $X$  be a graph,  $G \leq \text{Aut}(X)$ ,  $N \trianglelefteq G$  and  $X$  be  $N$ -regular covering of  $X_N$ . Then  $X$  is  $G$ -half-arc-transitive if and only if  $X_N$  is  $G/N$ -half-arc-transitive.*

*Proof.* Suppose that  $N \trianglelefteq G$  and  $X$  is  $G$ -half-arc-transitive. Since  $X$  is  $N$ -regular covering of  $X_N$ , it follows that  $K = N$  and  $G/N \leq \text{Aut}(X_N)$ , where  $K$  is the kernel of  $G$  acting on orbits of  $N$ . Let  $x^N, y^N$  be two arbitrary vertices of graph  $X_N$ . By our assumption there exists  $g \in G$  such that  $x^g = y$ . Now  $(x^N)^{Ng} = (x^g)^N = y^N$ . It implies that  $X_N$  is  $G/N$ -vertex-transitive. Now, suppose that  $\{x^N, y^N\}$  and  $\{u^N, v^N\}$  are two arbitrary edges of  $X_N$ . Without loss of generality, we may suppose that  $\{x, y\}$  and  $\{u, v\}$  are two edges of  $X$ . By our assumption there exists  $g \in G$  such that  $\{x, y\}^g = \{u, v\}$ . Then we may assume that  $x^g = u$  and  $y^g = v$ . Hence  $(x^N)^{Ng} = x^{Ng} = x^{gN} = u^N$  and  $(y^N)^{Ng} = y^{Ng} = y^{gN} = v^N$ . Then  $X_N$  is  $G/N$ -edge-transitive. Suppose to contrary that  $X_N$  is  $G/N$ -arc-transitive. Let  $(x, y)$  and  $(u, v)$  are two arcs of graph  $X$ . Now  $(x^N, y^N)$  and  $(u^N, v^N)$  are two arcs of graph  $X_N$ . By our assumption, there exists  $Ng \in G/N$  such that  $(x^N, y^N)^{Ng} = (u^N, v^N)$ . Therefore  $(x^N)^{Ng} = u^N$  and  $(y^N)^{Ng} = v^N$ . Thus  $x^{Ng} = u^N$  and  $y^{Ng} = v^N$ . Then  $x^g = u^n$  and  $y^g = v^{n'}$  for  $n, n' \in N$  and so  $(x, y)^g = (u^n, v^{n'})$ . There exists  $n'' \in N$  such that  $(u^n, v^{n'})^{n''} = (u, v)$ . Then  $(x, y)^{gn''} = (u^n, v^{n'})^{n''} = (u, v)$ . Therefore  $X$  is  $G$ -arc-transitive, a contradiction. Then  $X_N$  is  $G/N$ -half-arc-transitive.

Now suppose that  $X_N$  is  $G/N$ -half-arc-transitive. Thus  $G/N$  acts transitively on  $V(X_N)$ . Let  $u, v \in V(X)$  and  $u^N, v^N \in V(X_N)$ . Then there is  $Ng \in G/N$  such that  $(u^N)^{Ng} = v^N$  and hence, there is  $n' \in N$  such that  $u^g = v^{n'}$  and  $u^{g(n')^{-1}} = v$ . Then since  $g(n')^{-1} \in G$ , it implies that  $X$  is vertex-transitive. For any  $\{u, v\}, \{x, y\} \in E(X)$ , we have  $\{u^N, v^N\}, \{x^N, y^N\} \in E(X_N)$ . Since  $X_N$  is  $G/N$ -edge-transitive, we have  $Ng \in G/N$  such that  $\{u^N, v^N\}^{Ng} = \{x^N, y^N\}$  and  $\{(u^N)^{Ng}, (v^N)^{Ng}\} = \{x^N, y^N\}$ . Without loss of generality, we may suppose that  $(u^N)^{Ng} = (u)^{Ng} = x^N$  and  $(v^N)^{Ng} = (v)^{Ng} = y^N$ . There exists  $n', n'' \in N$  such that  $\{u, v\}^g = \{x^{n'}, y^{n''}\}$ . Also there exists  $n \in N$  such that  $\{x^{n'}, y^{n''}\}^n = \{x, y\}$ . Thus we may assume that  $\{u, v\}^{gn} = \{x, y\}$  and so  $X$  is  $G$ -edge-transitive. Similar to the previous, it can be shown that if  $X_N$  is not  $G/N$ -arc-transitive then  $X$  is not  $G$ -arc-transitive. Therefore  $X$  is  $G$ -half-arc-transitive. □

The following lemma is basic for the main result.

**Lemma 2.** *Let  $X$  be a half-arc-transitive graph,  $p$  is a prime and  $N \trianglelefteq \text{Aut}(X)$ , where  $N \cong \mathbb{Z}_p$ . If the quotient graph  $X_N$  is a Cayley graph and has the same valency with  $X$  then  $X$  is a  $N$ -regular covering of  $X_N$  and  $X$  is a Cayley graph.*

*Proof.* Let  $N$  be a normal subgroup of  $A := \text{Aut}(X)$  and  $X_N$  be the quotient graph of  $X$  with respect to the orbits of  $N$  on  $V(X)$ . Assume that  $K$  is the kernel of  $A$  acting on  $V(X_N)$ . The stabilizer  $K_v$  of  $v \in V(X)$  in  $K$  fixes the neighborhood of  $v$  in  $X$ . The connectivity of  $X$  implies  $K_v = 1$  for any  $v \in V(X)$  and hence  $N_v = 1$ . If  $N_{\{\alpha, \beta\}} \neq 1$  then  $N_{\{\alpha, \beta\}} = N$ , because  $N \cong \mathbb{Z}_p$ . Since  $X$  is connected, there is a  $\{\beta, \gamma\} \in E(X)$  where  $\beta, \gamma \in V(X)$ . Then we have  $g \in A$  such that  $\{\alpha, \beta\} = \{\beta, \gamma\}^g$  because  $X$  is

an edge-transitive graph. Hence  $N_{\{\alpha,\beta\}} = N_{\{\beta,\gamma\}} = g^{-1}N_{\{\beta,\gamma\}}g = N_{\{\beta,\gamma\}}$ . It is a contradiction and so  $N_{\{\alpha,\beta\}} = 1$ . Therefore  $X$  is a  $\mathbb{Z}_p$ -regular covering of  $X_N$ . Now we prove that  $X$  is a Cayley graph. Let  $X_N \cong \text{Cay}(G, S)$ ,  $X \cong X_N \times_{\xi} \mathbb{Z}_p$  where  $\xi$  is the  $T$ -reduced voltage assignment and  $\tilde{G}$  is a lift of  $G$  such that  $\tilde{\alpha}p = p\alpha$  where  $p : X \rightarrow X_N$  is regular covering projection,  $\alpha \in \text{Aut}(X_N)$  and  $\tilde{\alpha} \in A$ . For any  $(x, k), (y, k') \in V(X)$  where  $k, k' \in \mathbb{Z}_p$  and  $x, y \in V(X_N)$ , we have  $\alpha \in \text{Aut}(X_N)$  such that  $x^\alpha = y$ . For  $k'' \in \mathbb{Z}_p$ ,  $(x, k)^{\tilde{\alpha}p} = (z, k'')^p = z$  where  $(x, k)^{\tilde{\alpha}} = (z, k'')$ . Also  $(x, k)^{p\alpha} = x^\alpha = y$ . Then  $y = z$  and hence  $(y, k), (y, k'') \in p^{-1}(y)$ . Therefore  $\tilde{G}$  is transitive on  $V(X)$ . Now, we prove that  $\tilde{G}$  is semiregular. Suppose that  $(x, k)^{\tilde{\alpha}} = (x, k)$ . Now, since  $G$  is semiregular and  $\tilde{\alpha}p = p\alpha$ , it implies that  $x = (x, k)^{\tilde{\alpha}p} = (x, k)^{p\alpha} = x^\alpha$ . Then  $\alpha = 1$  and hence  $\tilde{\alpha}p = p$ . Therefore  $\tilde{\alpha} \in \text{CT}(p) = \mathbb{Z}_p$  and since  $\text{CT}(p)$  is semiregular, it follows that  $\tilde{\alpha} = 1$ . □

By [27], all tetravalent half-arc-transitive graphs of order  $12p$  where  $p \leq 53$  is a prime, are classified. Then in the following, we may assume that  $p > 53$ .

**Lemma 3.** *Let  $X$  be a tetravalent half-arc-transitive graph of order  $12p$ , where  $p$  is a prime. Then  $\text{Aut}(X)$  has a normal Sylow  $p$ -subgroup or  $X$  is  $\mathbb{Z}_3$ -regular cover of  $C(2; p, 2)$  or  $C_{2p}[2K_1]$ .*

*Proof.* Let  $X$  be a tetravalent half-arc-transitive graph of order  $12p$  where  $p$  is a prime. Let  $A := \text{Aut}(X)$ . Since the stabilizer  $A_v$  of  $v \in V(X)$  is a 2-group, we have  $|A| = 2^{m+2}.3.p$ , for some nonnegative integer  $m$ . Suppose to the contrary that  $A$  has no normal Sylow  $p$ -subgroups. Let  $N$  be a minimal normal subgroup of  $A$ . We claim that  $N$  is solvable. Otherwise, by Proposition 4 and since  $p > 53$ , we get a contradiction. Then  $N$  is solvable and hence it is an elementary abelian 2-,3- or  $p$ -group.

**Case I.**  $N$  is a 2-group.

Let  $X_N$  be the quotient graph of  $X$  corresponding to the orbits of  $N$  on  $V(X)$ . Then  $|V(X_N)| = 6p$  or  $3p$ .

**Subcase 1.**  $|V(X_N)| = 6p$ .

Since  $X$  is edge-transitive,  $X_N$  has valency 2 or 4. Suppose that  $X_N$  has valency 2. Then  $X \cong C_{6p}[2K_1]$ , which is arc-transitive. It is a contradiction. Assume now that  $X_N$  has valency 4. If  $X_N$  is half-arc-transitive then by [12, Theorem 4.1],  $|\text{Aut}(X_N)| = 2^2.3.p$ . Let  $K$  be the kernel of  $A$  acting on  $V(X_N)$ . Since  $K$  fixes each orbit of  $N$ , the stabilizer  $K_v = 1$  for any  $v \in V(X)$ . Then  $|N| = |K|$ . On the other hand  $A/K \leq \text{Aut}(X_N)$ . Since  $A/K$  acts transitively on  $V(X_N)$  and  $E(X_N)$ ,  $|A| = 24p$ . Then  $1 + np \mid 24$ . Since  $p > 53$  then  $P \trianglelefteq A$ , a contradiction. Now, suppose that  $X_N$  is arc-transitive. Let  $X_N$  has valency 4. By Proposition 5, if  $X_N$  is arc-regular then  $|\text{Aut}(X_N)| = 24p$ . By lemma 1,  $A/K$  is half-arc-transitive and hence  $|A| = 24p$ . Then  $P \trianglelefteq A$  because  $p > 53$ . It is a contradiction. If  $X_N$  do not be arc-regular then by Proposition 5,  $Y = X_N \cong C_{3p}[2K_1]$  and  $B = \text{Aut}(Y)$ .  $|B| = 2^{3p+1}.3.p$ . Assume that  $M$  is a minimal normal subgroup of  $B$ . By the same argument as in the first paragraph,  $M$  is solvable and hence it is an elementary abelian

2-,3- or  $p$ -group. First, assume that  $M$  is a 2-group and  $Y_M$  is the quotient graph of  $Y$  corresponding to the orbits of  $M$  on  $V(Y)$ . The quotient graph  $Y_M$  has order  $3p$  and valency 2 or 4. If  $Y_M$  has valency 4 then  $M_v = 1$  for  $v \in V(Y)$ . Assume that  $K_1$  be the kernel of  $B$  acting on  $V(Y_M)$ . Hence  $|K_1| = |M|$ . Thus  $B/K_1 \leq \text{Aut}(Y_M)$ . It is a contradiction because  $|\text{Aut}(Y_M)| = 12p$  by [31, Theorem 5]. If  $Y_M$  has valency 2 then  $Y_M \cong C_{3p}$  and  $\text{Aut}(Y_M) \cong D_{6p}$ . Since  $|K_1| \leq 2$ , we have  $|B| \leq 12p$ . We get a contradiction because  $p > 53$ . Now, suppose that  $M$  be a 3-group. Then  $|V(Y_M)| = 2p$ . Since  $M_v = 1$  for  $v \in V(Y)$  by using [16, Theorem 1.1(4)],  $Y_M$  has valency 4. By [8, Table 1],  $Y_M \cong G(2, p, r)$  or  $G(2p, r)$ . Then  $|K_1| = |M|$  and hence  $B/K_1 \leq \text{Aut}(Y_M)$ . It is a contradiction because  $|\text{Aut}(Y_M)| = 2^{p+1} \cdot p$  or  $8p$  and  $p > 53$ . Let  $M$  be a  $p$ -group. Then  $|Y_M| = 6$ . Since  $M_v \leq M$  we have  $|M_v| = 1$ . By [16, Theorem 1.1(4)],  $Y_M$  has valency 4. By [27],  $|\text{Aut}(Y_M)| = 48$ . Hence  $B/K_1 \leq \text{Aut}(Y_M)$ . It is a contradiction.

**Subcase 2.**  $|V(X_N)| = 3p$ .

Let  $|V(X_N)| = 3p$  and  $X_N$  has valency 2. Then  $X \cong C_{3p}[2K_1]$ . This leads to a contradiction. If  $X_N$  has valency 4 and it is half-arc-transitive then by [1, Theorem 2.5],  $|\text{Aut}(X_N)| = 6p$ . Since  $X_N$  is an edge-transitive graph,  $6p \mid |A/K| \mid 6p$ . Then  $|A| = 24p$  and hence  $P \trianglelefteq A$ . It is a contradiction. Suppose now that  $X_N$  is arc-transitive. By [31, Theorem 5],  $|\text{Aut}(X_N)| = 12p$ . Then with the same arguments as before, a contradiction can be obtained.

**Case II.**  $N$  is 3-group.

If  $|V(X_N)| = 4p$  and  $X_N$  has valency 2, then  $X_N \cong C_{4p}$  and hence  $\text{Aut}(X_N) \cong D_{8p}$ . Since  $K = K_v N$  for any  $v \in V(X)$  and  $K$  acts faithfully on  $V(X)$ , we have  $K \leq S_3$  and hence  $K_v \leq 2$ . Then  $|A| \mid 48p$ . Therefore  $P \trianglelefteq A$  according to assumption  $p > 53$ . This leads to a contradiction. Now let  $|V(X_N)| = 4p$  and  $X_N$  has valency 4. Then  $X_N$  is arc-transitive or half-arc-transitive. By [39, Table 1] and Proposition 2,  $X_N \cong C(2; p, 2)$ ,  $C_{2p}[2K_1]$ ,  $CA_{4p}^0$ ,  $CA_{4p}^1$  or  $X(4, p)$ . Let  $X_N \cong C(2; p, 2)$  or  $C_{2p}[2K_1]$ . Since  $X_N$  has valency 4,  $N$  acts semiregularly on  $V(X)$  and so  $X$  is a  $\mathbb{Z}_3$ -regular cover of  $C(2; p, 2)$  or  $C_{2p}[2K_1]$ . Assume that  $Y = X_N \cong CA_{4p}^0$  or  $CA_{4p}^1$  and  $B = \text{Aut}(Y)$ . Since  $|K| = |N|$ , we have  $A/K \leq B$  and hence  $|A| \leq 48p$ . Then  $P \trianglelefteq A$ . Suppose that  $Y = X_N \cong X(4, p)$  and  $B = \text{Aut}(Y)$ . Since  $Y$  is half-arc-transitive, we have  $|B| = 2^{m+2} \cdot p$ , for some nonnegative integer  $m$ . Let  $M$  be a minimal normal subgroup of  $B$ . Thus  $M$  is an elementary abelian 2- or  $p$ -group. First, assume that  $M$  be a  $p$ -group and  $Y_M$  be the quotient graph of  $Y$  corresponding to the orbits of  $M$  on  $V(Y)$ . Then  $|V(Y_M)| = 4$ . Since  $Y$  is an edge-transitive graph and  $M_v = 1$  for  $v \in V(Y)$ ,  $Y_M$  has valency 4, a contradiction. Suppose that  $M$  is a 2-group. Therefore  $|V(Y_M)| = 2p$  or  $p$  and  $Y_M$  has valency 2 or 4.

**Subcase 1.**  $|V(Y_M)| = 2p$ .

If  $Y_M$  has valency 2 then  $Y \cong C_{2p}[2K_1]$ , which is arc-transitive. Since  $Y$  is half-arc-transitive, we get a contradiction. Suppose now that  $Y_M$  has valency 4. By [8, Table 1],  $Y_M \cong G(2p, 4)$  or  $G(2, p, 2)$ . Assume that  $Y_M \cong G(2p, 4)$ . Since  $(K_1)_v = 1$ ,  $|B/K_1| \leq 8p$  and hence  $|A| \leq 48p$ . It is a contradiction because  $p > 53$ . Suppose that  $Y_M \cong G(2, p, 2)$ . Let  $Z = Y_M \cong G(2, p, 2)$  and  $C = \text{Aut}(Z)$ . Let  $H$  be a minimal


normal subgroup of  $C$  and let  $Z_H$  be the quotient graph of  $Z$  with respect to the orbits of  $H$ . Since  $|C| = 2^{p+1} \cdot p$ ,  $H$  is 2- or  $p$ -group. Assume that  $H$  is a 2-group. Thus  $|Z_H| = p$  and  $Z_H$  has valency 2 or 4. By [6, Theorem 3],  $|\text{Aut}(Z_H)| = 2p$  or  $4p$ . Assume that  $K_1$  be the kernel of  $C$  acting on  $V(Z_H)$ . If  $Z_H$  has valency 4 then  $|K_1| = |H| = 2$  because  $|(K_1)_v| = 1$ . Then  $C/K_1 \leq 16p$  and hence  $2^{p+1} \leq 8p$ . We get a contradiction because  $p > 53$ . If  $Z_H$  has valency 2 then  $|K_1| \leq 8$  because  $|(K_1)_v| \leq 2$ . Thus  $C/K_1 \leq 16p$  and hence  $2^{p+1} \leq 8p$ , a contradiction can be obtained. Now, suppose that  $H$  is a  $p$ -group. Then  $|Z_H| = 2$  with valency 2, 4, a contradiction.

**Subcase 2:**  $|V(Y_M)| = p$ .

If  $Y_M$  has valency 4 then by lemma 2,  $Y$  is  $\mathbb{Z}_2$ -regular cover of  $Y_M$  and  $Y$  is a Cayley graph. But by [15],  $X(4, p)$  is not a Cayley graph, a contradiction. Suppose that  $Y_M$  has valency 2 and hence  $Y_M \cong C_p$ . Assume that  $K_1$  is the kernel of  $B$  acting on  $V(Y_M)$  and  $(K_1)_v = 1$ . Then  $B/K_1 \leq \text{Aut}(Y_M)$  and so  $|B| \leq 8p$ . Therefore  $|A| \leq 24p$  and hence  $P \trianglelefteq A$  because  $p > 53$ . Then  $(K_1)_v \neq 1$ . Let  $V(Y_M) = \{\Omega_0, \Omega_1, \Omega_2, \dots, \Omega_{p-1}\}$ . The subgraph induced by any two adjacent orbits is either a cycle of length 8 or a union of two cycles of length 4. Suppose that  $\langle \Omega_i \cup \Omega_{i+1} \rangle$  is an 8-cycle. Thus  $K_1$  acts faithfully on each  $\Omega_i$  and hence  $(K_1)_v \cong \mathbb{Z}_2$ . It implies that  $|K_1| = 8$ . Since  $M$  is transitive on each  $\Omega_i$  and  $(K_1)_v > 1$ , all edges in the induced subgraph  $\langle \Omega_i \cup \Omega_{i+1} \rangle$  have the same direction either from  $\Omega_i$  to  $\Omega_{i+1}$  or from  $\Omega_{i+1}$  to  $\Omega_i$  in the oriented graph  $D_B(Y)$ . It follows that  $B/K_1 \cong \mathbb{Z}_p$  and  $|B| \leq 8p$ . Therefore  $|A| \leq 24p$  and hence  $P \trianglelefteq A$  because  $p > 53$ . Assume that  $\langle \Omega_i \cup \Omega_{i+1} \rangle$  is a union of two 4-cycles. Let  $\Omega_i = \{u_i^0, u_i^1, u_i^2, u_i^3\}$  for any  $i$  in  $\mathbb{Z}_p$ . Then  $B$  has an automorphism  $\alpha$  of order  $p$  such that for any  $i$  in  $\mathbb{Z}_p$ ,  $\Omega_i^\alpha = \Omega_{i+1}$ . Let  $(u_i^j)^\alpha = u_{i+1}^j$  for  $i$  in  $\mathbb{Z}_p$  and  $j$  in  $\mathbb{Z}_4$ . Consider a 4-cycle  $C$  in the induced subgraph  $\langle \Omega_0 \cup \Omega_1 \rangle$  and let  $n$  be the number of edges of  $C$  which are in some orbit of  $\alpha$ . Then  $n = 0, 1$ , or 2. Consequently, the induced subgraph  $\langle \Omega_0 \cup \Omega_1 \rangle$  is one of the of the following three cases.

In the Case 1,  $Y$  is disconnected, a contradiction. In the Case 2,  $Y \cong C_{2p}[2K_1]$ . We get a contradiction because  $Y \cong X(4, p)$ . In the Case 3,  $Y \cong C(2; p, 2)$  that is arc-transitive. It is a contradiction because  $X(4, p)$  is a half-arc-transitive graph.

**Case III.**  $N$  is  $p$ -group.



**Figure 3.** The induced subgraph  $\langle \Omega_0 \cup \Omega_1 \rangle$

If  $|N| = p$  then  $N$  is a normal Sylow  $p$ -subgroup of  $A$  as claimed. □

**Theorem 1.** *Let  $X$  be a connected tetravalent half-arc-transitive graph of order  $12p$ , where  $p > 53$  is a prime. Then one of the following statements holds:*

- (1)  $X$  is  $\mathbb{Z}_3$ -regular cover of  $C(2; p, 2)$ .
- (2)  $X$  is  $\mathbb{Z}_3$ -regular cover of  $C_{2p}[2K_1]$  and in this case  $X$  is a Cayley graph.
- (3)  $X \cong X(r; 12, p)$  such that  $r \in \mathbb{Z}_p^*$  satisfying  $r^{12} = \pm 1$ .
- (4)  $X$  is  $\mathbb{Z}_p$ -regular cover of  $W(6, 2)$  or  $R_6(5, 4)$  and in this case  $X$  is a Cayley graph.

*Proof.* Let  $X$  be a tetravalent half-arc-transitive graph of order  $12p$  and hence  $|A| = 2^{m+2} \cdot 3 \cdot p$  for some integer  $m \geq 0$ . By Lemma 3, either  $P \trianglelefteq A$  or  $X$  is a  $\mathbb{Z}_3$ -regular cover of  $C(2; p, 2)$  or  $C_{2p}[2K_1]$ . If  $X$  is  $\mathbb{Z}_3$ -regular cover of  $C(2; p, 2)$  then we have Case 1. Also, if  $X$  is  $\mathbb{Z}_3$ -regular cover of  $C_{2p}[2K_1]$  then by Lemma 2,  $X$  is a Cayley graph and we have Case 2. Now, suppose that  $P \trianglelefteq A$ . Let  $X_P$  be the quotient graph of  $X$  corresponding to the orbits of  $P$ . Assume that  $K$  is the kernel of  $A$  acting on  $V(X_P)$ . Then  $V(X_P) = 12$  and  $X_P$  has valency 2 or 4. If  $X_P$  has valency 2 then  $X_P \cong C_{12}$  and hence  $\text{Aut}(X_P) \cong D_{24}$ . By Proposition 3,  $A_v \cong \mathbb{Z}_2$  and hence  $|A| = 24p$ . The attachment number of  $X$  is equal to its radius. So  $X$  is a tetravalent tightly attached half-arc-transitive graph of odd radius  $p$ . By Proposition 1,  $X \cong X(r; 12, p)$  where  $r \in \mathbb{Z}_p^*$  and  $r^{12} = \pm 1$ , which is Case 3. Assume that  $X_P$  has valency 4 and  $X_P$  is arc-transitive or half-arc-transitive. There is no half-arc-transitive graph of order 12. Suppose that  $X_P$  is an arc-transitive graph. By [27],  $W(6, 2)$  and  $R_6(5, 4)$  are the only two arc-transitive graphs of order 12. These graphs are Cayley graphs by [3]. Since  $P$  acts semiregular on  $V(X)$  and  $E(X)$ , by Lemma 2,  $X$  is a  $\mathbb{Z}_p$ -regular cover of  $X_P$  and  $X$  is a Cayley graph, which is Case 4.  $\square$

**Conflict of interest.** The authors declare that they have no conflict of interest.

**Data Availability.** Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

## References

- [1] B. Alspach and M.Y. Xu, *1/2-transitive graphs of order  $3p$* , J. Algebraic Combin. **3** (1994), no. 4, 347–355  
<https://doi.org/10.1023/A:1022466626755>.
- [2] I. Antončič and P. Šparl, *Classification of quartic half-arc-transitive weak metacirculants of girth at most 4*, Discrete Math. **339** (2016), no. 2, 931–945  
<https://doi.org/10.1016/j.disc.2015.10.015>.
- [3] R.A. Beezer, *Sage for linear algebra a supplement to a first course in linear algebra*, Sage web site <http://www.sagemath.org> (2011).
- [4] K. Berčič and M. Ghasemi, *Tetravalent arc-transitive graphs of order twice a product of two primes*, Discrete Math. **312** (2012), no. 24, 3643–3648  
<https://doi.org/10.1016/j.disc.2012.08.018>.

- [5] I.Z. Bouwer, *Vertex and edge transitive, but not 1-transitive, graphs*, Can. Math. Bull. **13** (1970), no. 2, 231–237  
<https://doi.org/10.4153/CMB-1970-047-8>.
- [6] C.Y. Chao, *On the classification of symmetric graphs with a prime number of vertices*, Trans. Amer. Math. Soc. **158** (1971), no. 1, 247–256  
<https://doi.org/10.2307/1995785>.
- [7] H. Cheng and L. Cui, *Tetravalent half-arc-transitive graphs of order  $p^5$* , Appl. Math. Comput. **332** (2018), 506–518  
<https://doi.org/10.1016/j.amc.2018.03.076>.
- [8] Y. Cheng and J. Oxley, *On weakly symmetric graphs of order twice a prime*, J. Combin. Theory, Ser. B **42** (1987), no. 2, 196–211  
[https://doi.org/10.1016/0095-8956\(87\)90040-2](https://doi.org/10.1016/0095-8956(87)90040-2).
- [9] M.D.E. Conder and A. Žitnik, *Half-arc-transitive graphs of arbitrary even valency greater than 2*, European J. Combin. **54** (2016), 177–186  
<https://doi.org/10.1016/j.ejc.2015.12.011>.
- [10] L. Cui and J.X. Zhou, *A classification of tetravalent half-arc-transitive metacirculants of 2-power orders*, Appl. Math. Comput. **392** (2021), Article ID: 125755  
<https://doi.org/10.1016/j.amc.2020.125755>.
- [11] S.F. Du and M.Y. Xu, *Vertex-primitive 1/2-arc-transitive graphs of smallest order*, Commun. Algebra **27** (1999), 163–171.
- [12] Y.Q. Feng, J.H. Kwak, X. Wang, and J.X. Zhou, *Tetravalent half-arc-transitive graphs of order  $2pq$* , J. Algebraic Combin. **33** (2011), no. 4, 543–553  
<https://doi.org/10.1007/s10801-010-0257-1>.
- [13] Y.Q. Feng, J.H. Kwak, M.Y. Xu, and J.X. Zhou, *Tetravalent half-arc-transitive graphs of order  $p^4$* , European J. Combin. **29** (2008), no. 3, 555–567  
<https://doi.org/10.1016/j.ejc.2007.05.004>.
- [14] Y.Q. Feng, J.H. Kwak, and C. Zhou, *Constructing even radius tightly attached half-arc-transitive graphs of valency four*, J. Algebraic Combin. **26** (2007), no. 4, 431–451  
<https://doi.org/10.1007/s10801-007-0064-5>.
- [15] Y.Q. Feng, K. Wang, and C. Zhou, *Tetravalent half-transitive graphs of order  $4p$* , European J. Combin. **28** (2007), no. 3, 726–733  
<https://doi.org/10.1016/j.ejc.2006.01.002>.
- [16] A. Gardiner and C.E. Praeger, *On 4-valent symmetric graphs*, European J. Combin. **15** (1994), no. 4, 375–381  
<https://doi.org/10.1006/eujc.1994.1041>.
- [17] D. Gorenstein, *Finite simple groups*, Plenum, New York, 1982.
- [18] J.L. Gross and T.W. Tucker, *Generating all graph coverings by permutation voltage assignments*, Discrete Math. **18** (1977), no. 3, 273–283  
[https://doi.org/10.1016/0012-365X\(77\)90131-5](https://doi.org/10.1016/0012-365X(77)90131-5).
- [19] D.F. Holt, *A graph which is edge transitive but not arc transitive*, J. Graph Theory **5** (1981), no. 2, 201–204  
<http://doi.org/10.1002/jgt.3190050210>.
- [20] A. Hujdurović, K. Kutnar, and D. Marušič, *Half-arc-transitive group actions with*

- a small number of alternets*, J. Combin. Theory, Ser. A **124** (2014), no. 1, 114–129  
<http://doi.org/10.1016/j.jcta.2014.01.005>.
- [21] K. Kutnar, D. Marušič, and P. Šparl, *An infinite family of half-arc-transitive graphs with universal reachability relation*, European J. Combin. **31** (2010), no. 7, 1725–1734  
<https://doi.org/10.1016/j.ejc.2010.03.006>.
- [22] K. Kutnar, D. Marušič, P. Šparl, R.J. Wang, and M.Y. Xu, *Classification of half-arc-transitive graphs of order  $4p$* , European J. Combin. **34** (2013), no. 7, 1158–1176  
<https://doi.org/10.1016/j.ejc.2013.04.004>.
- [23] C.H. Li and H.S. Sim, *On half-transitive metacirculant graphs of prime-power order*, J. Combin. Theory, Ser. B **81** (2001), no. 1, 45–57  
<https://doi.org/10.1006/jctb.2000.1992>.
- [24] H. Liu, B. Lou, and B. Ling, *Tetravalent half-arc-transitive graphs of order  $p^2q^2$* , Czechoslovak Math. J. **69** (2019), no. 2, 391–401  
<https://doi.org/10.21136/CMJ.2019.0335-17>.
- [25] D. Marušič, *Half-transitive group actions on finite graphs of valency 4*, J. Combin. Theory, Ser. B **73** (1998), no. 1, 41–76  
<https://doi.org/10.1006/jctb.1997.1807>.
- [26] D. Marušič and C.E. Praeger, *Tetravalent graphs admitting half-transitive group actions: alternating cycles*, J. Combin. Theory, Ser. B **75** (1999), no. 2, 188–205  
<https://doi.org/10.1006/jctb.1998.1875>.
- [27] P. Potočník and S. Wilson, *A census of edge-transitive tetravalent graphs*, Available at <http://jan.ucc.nau.edu/swilson/C4fullSite/index.html>.
- [28] C.E. Praeger and M.Y. Xu, *A characterisation of a class of symmetric graphs of twice prime valency*, European J. Combin. **10** (1986), 91–102  
[https://doi.org/10.1016/S0195-6698\(89\)80037-X](https://doi.org/10.1016/S0195-6698(89)80037-X).
- [29] W.T. Tutte, *Connectivity in Graphs*, University of Toronto Press, 1966.
- [30] R.J. Wang, *Half-transitive graphs of order a product of two distinct primes*, Commun. Algebra **22** (1994), no. 3, 915–927  
<https://doi.org/10.1080/00927879408824885>.
- [31] R.J. Wang and M.Y. Xu, *A classification of symmetric graphs of order  $3p$* , J. Combin. Theory, Ser. B **58** (1993), no. 2, 197–216  
<https://doi.org/10.1006/jctb.1993.1037>.
- [32] X. Wang, Y. Feng, J. Zhou, J. Wang, and Q. Ma, *Tetravalent half-arc-transitive graphs of order a product of three primes*, Discrete Math. **339** (2016), no. 5, 1566–1573  
<https://doi.org/10.1016/j.disc.2015.12.022>.
- [33] X. Wang and Y.Q. Feng, *Half-arc-transitive graphs of order  $4p$  of valency twice a prime*, Ars Math. Contemp. **3** (2010), no. 2, 151–163  
<http://doi.org/10.26493/1855-3974.125.164>.
- [34] \_\_\_\_\_, *There exists no tetravalent half-arc-transitive graph of order  $2p^2$* , Discrete Math. **310** (2010), no. 12, 1721–1724

- <https://doi.org/10.1016/j.disc.2009.11.020>.
- [35] X. Wang, J. Wang, and Y. Liu, *Tetravalent half-arc-transitive graphs of order  $8p$* , J. Algebraic Combin. **51** (2019), 237–246  
<https://doi.org/10.1007/s10801-019-00873-y>.
- [36] Y. Wang and Y.Q. Feng, *Half-arc-transitive graphs of prime-cube order of small valencies*, Ars Math. Contemp. **13** (2017), no. 2, 343–35  
<http://doi.org/10.26493/1855-3974.964.594>.
- [37] M.Y. Xu, *Half-transitive graphs of prime-cube order*, J. Algebraic Combin. **1** (1992), no. 3, 275–282  
<https://doi.org/10.1023/A:1022440002282>.
- [38] M.M. Zhang and J.X. Zhou, *Tetravalent half-arc-transitive bi- $p$ -metacirculants*, J. Graph Theory **92** (2019), no. 1, 19–38  
<https://doi.org/10.1002/jgt.22438>.
- [39] J.X. Zhou, *Tetravalent  $s$ -transitive graphs of order  $4p$* , Discrete Math. **309** (2009), no. 20, 6081–6086  
<https://doi.org/10.1016/j.disc.2009.05.014>.
- [40] J.X. Zhou and Y.Q. Feng, *Tetravalent one-regular graphs of order  $2pq$* , J. Algebraic Combin. **29** (2009), 457–471  
<https://doi.org/10.1007/s10801-008-0146-z>.