

## Leech graphs

Seena Varghese<sup>1,\*</sup>, Aparna Lakshmanan Savithri<sup>2</sup> and S. Arumugam<sup>3</sup>

<sup>1</sup>Department of Mathematics, Federal Institute of Science and Technology,  
Angamaly - 683577, Kerala, India  
\*graceseenaprince@gmail.com

<sup>2</sup>Department of Mathematics, Cochin University of Science and Technology,  
Cochin-22, Kerala, India  
aparnals@cusat.ac.in

<sup>3</sup>National Centre for Advanced Research in Discrete Mathematics, Kalasalingam University  
Anand Nagar, Krishnankoil-626 126, Tamil Nadu, India  
s.arumugam.klu@gmail.com

*Received: 31 March 2022; Accepted: 30 September 2022*  
*Published Online: 3 October 2022*

**Abstract:** Let  $t_p(G)$  denote the number of paths in a graph  $G$  and let  $f : E \rightarrow \mathbb{Z}^+$  be an edge labeling of  $G$ . The weight of a path  $P$  is the sum of the labels assigned to the edges of  $P$ . If the set of weights of the paths in  $G$  is  $\{1, 2, 3, \dots, t_p(G)\}$ , then  $f$  is called a Leech labeling of  $G$  and a graph which admits a Leech labeling is called a Leech graph. In this paper, we prove that the complete bipartite graphs  $K_{2,n}$  and  $K_{3,n}$  are not Leech graphs and determine the maximum possible value that can be given to an edge in the Leech labeling of a cycle.

**Keywords:** Leech labeling, Leech tree, Leech graph

**AMS Subject classification:** 05C78

### 1. Introduction

By a graph  $G = (V, E)$  we mean a finite undirected graph with neither loops nor multiple edges. The order  $|V|$  and the size  $|E|$  are denoted by  $n$  and  $m$  respectively. For graph theoretic terminology we refer to Chartrand and Lesniak [2].

Let  $f : E \rightarrow \mathbb{Z}^+$  be an edge labeling of  $G$ . The weight of a path  $P$  in  $G$  is the sum of the labels of the edges of  $P$  and is denoted by  $w(P)$ . Leech [5] introduced the concept of a Leech tree, while considering a problem in electrical engineering, where edge labels represent electrical resistance. Let  $T$  be a tree of order  $n$ . An edge labeling

---

\* Corresponding Author

$f : E \rightarrow \mathbb{Z}^+$  is called a Leech labeling if the weights of the  $\binom{n}{2}$  paths in  $T$  are exactly  $1, 2, \dots, \binom{n}{2}$ . A tree which admits a Leech labeling is called a Leech tree. Since each edge label is the weight of a path of length one, it follows that  $f$  is an injection and 1,2 are edge labels for all  $n \geq 3$ . Leech found five Leech trees which are given in Figure 1 and these are the only known Leech trees.




Figure 1. Leech trees

Taylor [8] proved that if  $T$  is a Leech tree of order  $n$ , then  $n = k^2$  or  $k^2 + 2$  for some integer  $k$ . Since then it has been proved by several authors ([1, 7, 9]) that no Leech trees of order 9, 11 or 16 exist, leaving  $n = 18$  as the smallest open case. In [13] and [10], it is shown that bistars, tristars and a subclass of trees of diameter  $n - 2$  are non-Leech trees. Some variations of Leech trees such as modular Leech trees ([3, 4]), minimal distinct distance trees [1] and leaf-Leech trees [6] have been investigated by several authors. A parameter called Leech index was introduced in [11], which measures how close a tree is towards being a Leech tree.

The total number of paths in a graph  $G$  is called the path number of  $G$  and is denoted by  $t_p(G)$ . Let  $f : E \rightarrow \mathbb{Z}^+$  be an edge labeling of  $G$ . The weight of a path  $P$  in  $G$  is the sum of the labels of the edges of  $P$  and is denoted by  $w(P)$ . If the set of weights of the paths in  $G$  is  $\{1, 2, 3, \dots, t_p(G)\}$ , then  $f$  is called a Leech labeling of  $G$  and a graph that admits a Leech labeling is called a Leech graph [12].

Let  $f$  be an edge labeling of a graph  $G$  such that both  $f$  and the weight function  $w$  on the set of all paths of  $G$  are both injective. Let  $S$  be the set of all path weights. Let  $k_f$  be the positive integer such that  $\{1, 2, 3, \dots, k_f\} \subseteq S$  and  $k_f + 1 \notin S$ . Let  $k(G) = \max k_f$ , where the maximum is taken over all such edge labelings  $f$ . Then  $k(G)$  is called the Leech index of the graph  $G$ .

In [12], it has been proved that cycles of order at most 6 are Leech graphs, whereas complete graphs of order 4, 5 and 6 are non-Leech graphs. The case  $n \geq 7$  is left as an open problem for both cycles and complete graphs. It is a simple observation that  $K_4 - \{e\}$  and  $P_5$  are non-Leech graphs of smallest order and smallest size. Since  $C_6$  is a Leech graph and  $P_5$  is not a Leech graph, it follows that the property of being a Leech graph is not hereditary and hence does not admit a forbidden subgraph characterization.

In this paper, we prove that  $K_{2,n}$  and  $K_{3,n}$ , for  $n > 2$  are non-Leech graphs and determine the Leech index of these graphs. We also prove some properties of Leech cycles.

## 2. Complete Bipartite Graphs

Throughout this section, let  $X = \{u_1, u_2, \dots, u_m\}$  and  $Y = \{v_1, v_2, \dots, v_n\}$  be the bipartition of  $K_{m,n}$ . If  $P_1$  and  $P_2$  are two paths having a common end vertex  $v$  and  $V(P_1) \cap V(P_2) = \{v\}$ , then the path obtained by concatenation of  $P_1$  and  $P_2$  is denoted by  $P_1 \circ P_2$ .

**Lemma 1.** *Let  $f$  be a Leech labeling of  $K_{m,n}$ ,  $m, n > 2$ . Let  $P_1$  be a  $x - y$  path,  $P_2$  be a  $r - s$  path,  $V(P_1) \cap V(P_2) = \phi$ ,  $w(P_1) = k$ ,  $w(P_2) = l$  and  $xr \in E(K_{m,n})$ . Then  $w(P) \neq k + l$  for any path  $P$  with  $x$  as origin and  $r \notin V(P)$ .*

*Proof.* Suppose there exists a path  $P$  with  $x$  as origin,  $r \notin V(P)$  and  $w(P) = k + l$ . Then the two paths  $P \circ (x, r)$  and  $P_1 \circ (x, r) \circ P_2$  have the weight  $k + l + f(xr)$ , which is a contradiction.  $\square$

**Corollary 1.** *Let  $f$  be a Leech labeling of  $K_{m,n}$ ,  $m, n > 2$ . Let  $e_1$  and  $e_2$  be two nonadjacent edges,  $f(e_1) = k$  and  $f(e_2) = l$ . Then  $f(e) \neq k + l$  for any edge  $e$  adjacent to  $e_1$  or  $e_2$ .*

**Corollary 2.** *Let  $f$  be a Leech labeling of  $K_{3,n}$  and let  $n > 3$ . Let  $P_1$  and  $P_2$  be two vertex disjoint paths such that the end vertices of  $P_1$  and  $P_2$  cover all the vertices of  $X = \{u_1, u_2, u_3\}$ . Let  $w(P_1) + w(P_2) = c$ . Then  $f(e) \neq c$  for any edge  $e$ .*

*Proof.* Suppose there exists an edge  $e$  with  $f(e) = c$ . Since the end vertices of  $P_1$  and  $P_2$  cover  $X$ , we may assume without loss of generality that  $u_1$  is an end vertex of  $P_1$  and  $e$ . We claim that no vertex of the other partite set  $Y = \{v_1, v_2, \dots, v_n\}$  is an end vertex of  $P_2$ . Suppose  $v_1$  is an end vertex of  $P_2$ . If  $e = u_1v_i$  where  $i \neq 1$ , then  $w(P_1 \circ (u_1, v_1) \circ P_2) = w((v_i, u_1, v_1)) = c + f(u_1v_1)$ , which is a contradiction. Hence  $e = u_1v_1$  and the other end vertex of  $P_2$  is  $u_2$  or  $u_3$ . Let  $u_2$  be the other end vertex of  $P_2$ . Since  $n > 3$ , there exists  $v_i$  in  $Y$  such that  $v_i \notin (V(P_1) \cup V(P_2))$ . Now,  $w(P_1 \circ (u_1, v_i, u_2) \circ P_2) = w((v_1, u_1, v_i, u_2)) = c + f(u_1v_i) + f(v_iu_2)$ , which is a contradiction. Thus no vertex of  $Y$  is an end vertex of  $P_2$  and hence  $P_2$  is a  $u_2$ - $u_3$  path. Clearly,  $P_1$  has length 1 and  $P_2$  has length 2. Let  $P_1 = (u_1, v_1)$ ,  $P_2 = (u_2, v_2, u_3)$  and  $e = u_1v_i$  where  $i \neq 1$ . Since  $n > 3$ , there exists  $v_j \in Y$  such that  $j \notin \{1, 2, i\}$ . Then  $w(P_1 \circ (u_1, v_j, u_2) \circ P_2) = w((v_i, u_1, v_j, u_2)) = c + f(u_1v_j) + f(v_ju_2)$ , which is a contradiction. Hence  $f(e) \neq c$  for any edge  $e$ .  $\square$

**Corollary 3.** *Let  $f$  be a Leech labeling of  $K_{3,n}$ ,  $n > 3$ . Let  $P_1$  and  $P_2$  be two vertex disjoint paths and let  $P_3$  and  $P_4$  be another pair of vertex disjoint paths such that  $w(P_1) + w(P_2) = w(P_3) + w(P_4) = c$  and the end vertices of  $P_1, P_2, P_3, P_4$  covers all the vertices of  $X = \{u_1, u_2, u_3\}$ . Then  $f(e) \neq c$  for any edge.*

*Proof.* Suppose there exists an edge  $e$  with  $f(e) = c$ . Since the end vertices of  $P_1, P_2, P_3$  and  $P_4$  cover  $X$ , we may assume without loss of generality that  $u_1$  is an end vertex of  $P_1$  and  $e$ . Proceeding as in Corollary 2, the proof follows.  $\square$

**Lemma 2.** *Let  $f$  be a Leech labeling of  $K_{2,n}$ ,  $n > 2$ . Let  $f(u_1v_i) + f(u_2v_j) = c$  where  $i \neq j$ . Then  $f(e) \neq c$  for any edge  $e$ .*

*Proof.* Suppose  $f(e) = c$  for some edge  $e$ . Since  $e$  is incident with  $u_1$  or  $u_2$ , we may assume that  $e = u_1v_k$  where  $k \neq i$ . If  $k \neq j$ , then  $w((v_k, u_1, v_j)) = w((v_i, u_1, v_j, u_2)) = c + f(u_1v_j)$ . If  $k = j$ , then for any  $r \neq i, j$ ,  $w((v_i, u_1, v_r, u_2, v_j)) = w((v_j, u_1, v_r, u_2)) = c + f(u_1v_r) + f(u_2v_r)$ . Hence the result follows.  $\square$

We now proceed to prove that  $K_{2,n}$  and  $K_{3,n}$  are not Leech graphs for all  $n \geq 3$ . Throughout the proof  $S$  denotes the set of all path weights at each stage. For any positive integer  $r$ , the set  $\{1, 2, \dots, r\}$  is denoted by  $[r]$ .

**Theorem 1.** *The complete bipartite graph  $K_{2,n}$ ,  $n > 2$  is not a Leech graph.*

*Proof.* Suppose  $K_{2,n}$  is a Leech graph with Leech labeling  $f$ . It follows from Lemma 2 that the edges with labels 1 and 2 are adjacent. Suppose  $f(u_1v_1) = 1$  and  $f(u_2v_1) = 2$ . Then either  $f(u_1v_2) = 4$  or  $f(u_2v_2) = 4$ . If  $f(u_1v_2) = 4$ , then  $[5] \subseteq S$  and there cannot be a path of weight 6. Similarly, if  $f(u_2v_2) = 4$ , then there cannot be a path of weight 5, which is a contradiction. Hence  $f(u_1v_1) = 1, f(u_1v_2) = 2$  and  $[3] \subseteq S$ . If 4 is assigned to an edge not adjacent to  $u_1v_1, u_1v_2$ , then it follows from Lemma 2 that the path weights 5 or 6 cannot be obtained. Hence let  $f(u_1v_3) = 4$ , so that  $[6] \subseteq S$ . Again it follows from Lemma 2 that if 7 is assigned to an edge not adjacent to  $u_1v_1, u_1v_2$ , then the path weight 8 or 9 cannot be obtained. Hence  $f(u_1v_4) = 7$ , so that  $[9] \cup \{11\} \subseteq S$ . Now let  $f(e) = 10$ . Since  $11 \in S$ ,  $e$  is not adjacent to  $u_1v_1$ . If  $e$  is not adjacent to  $u_1v_2$ , then by Lemma 2, the path weight 12 cannot be obtained. Hence  $e = u_2v_2$ . Now  $u_2v_2$  and  $u_1v_3$  are nonadjacent and by Lemma 2 the path weight 14 cannot be obtained. Hence  $K_{2,n}$  is not a Leech graph.  $\square$

**Corollary 4.** *The Leech index of  $K_{2,n}$  is  $k(K_{2,n}) = 13$ , for  $n \geq 4$ . When  $n = 3$ ,  $k(K_{2,3}) = 8$  and when  $n = 2$ ,  $K_{2,2} = C_4$  which is a Leech graph.*

**Theorem 2.** *The complete bipartite graph  $K_{3,n}$ ,  $n \geq 3$  is not a Leech graph.*

*Proof.* Suppose  $K_{3,n}$  is a Leech graph with Leech labeling  $f$ . Let  $f(e_1) = 1$  and  $f(e_2) = 2$ . Suppose  $e_1$  and  $e_2$  are nonadjacent. Then by Corollary 1, the edge  $e_3$  with  $f(e_3) = 3$  is nonadjacent to  $e_1$  and  $e_2$  and the edge  $e_4$  with  $f(e_4) = 4$  is nonadjacent to  $e_1$  and  $e_3$ . Thus 1 and 4 are assigned to a pair of nonadjacent edges and 2 and 3 are assigned to a pair of nonadjacent edges. Hence path weight 5 cannot be obtained. Thus  $e_1$  and  $e_2$  are adjacent. We consider two cases.

**Case 1.**  $e_1 = u_1v_1$  and  $e_2 = u_1v_2$ .

Hence  $[3] \subseteq S$ . Let  $f(e_3) = 4$ . If  $e_3$  is nonadjacent to  $e_1$  and  $e_2$ , then by Corollary 1, the label 5 must be assigned to an edge not adjacent to  $e_1$  and  $e_2$ . Thus 1 and 5 are assigned to two nonadjacent edges and 2 and 4 are assigned to two nonadjacent edges. Hence the path weight 6 cannot be obtained.

If  $e_3$  is nonadjacent to  $e_1$  and adjacent to  $e_2$ , then 5 must be assigned to an edge independent to  $e_1$  and  $e_3$ , say,  $f(u_3v_3) = 5$ . Then path weight  $6 + f(u_1v_3)$  repeats.

Now suppose  $e_3$  is adjacent to  $e_1$  and not adjacent to  $e_2$ . Let  $e_3 = u_2v_1$ . Then by Corollary 1, the label 6 must be assigned to an edge not adjacent to both  $e_2 = u_1v_2$  and  $e_3 = u_2v_1$ . Hence  $[7] \subseteq S$ . By Corollary 1 the label 8 must be assigned to an edge not adjacent to the edges with labels 2 and 6. Thus 1 and 8 are assigned to two nonadjacent edges and 3 and 6 are path weights of two vertex disjoint paths. Hence, by Corollary 3, path weight 9 cannot be obtained.

Therefore  $e_3$  is adjacent to both  $e_1$  and  $e_2$ . Let  $e_3 = u_1v_3$ . Hence  $[6] \subseteq S$ . Now let  $f(e_4) = 7$ . If  $e_4$  is not adjacent to  $e_1$ , then by Corollary 1, the label 8 must be assigned to an edge  $e$  not adjacent to  $e_1$  and  $e_4$ . Now if  $e_4$  is not adjacent to  $e_2$ , then  $f(e_2) + f(e_4) = f(e_1) + f(e) = 9$  and hence by Corollary 3, the path weight 9 cannot be obtained. If  $e_4$  is adjacent to  $e_2$ , then 8 must be assigned to an edge independent to  $e_1$  and  $e_4$ , say,  $f(u_3v_j) = 8, j \neq 1, 2$  and then  $9 + f(v_ju_1)$  repeats. Hence  $e_4$  is adjacent to  $e_1$ . Now if  $e_4$  is not adjacent to  $e_2$ , then by Corollary 1, the label 9 must be assigned to an edge  $e$  not adjacent to  $e_2$  and  $e_4$ . Hence  $f(e_3) + f(e_4) = f(e) + f(e_2) = 11$  and by Corollary 3, path weight 11 cannot be obtained. Thus  $e_4$  is adjacent to  $e_2$ . Therefore  $e_4 = u_1v_4$  and so  $[9] \cup \{11\} \subseteq S$ . Now let  $f(e_5) = 10$ .

Since  $11 \in S$ ,  $e_5$  is not adjacent to  $e_1$ . If  $e_5$  is adjacent to  $e_2$ , then  $[13] \cup \{16, 19\} \subseteq S$ . By Corollary 1, the label 14 must be assigned to an edge  $e$  not adjacent to  $e_3$  and  $e_5$ . If  $e$  is adjacent to  $e_1$ , then path weight 19 repeats. If  $e$  is non adjacent to  $e_1$ , then  $f(e_1) + f(e) = f(e_5) + w(v_1, u_1, u_3) = 15$ , and by Corollary 3, the path weight 15 cannot be obtained. Therefore  $e_5$  is not adjacent to  $e_2$ . Hence let  $e_5 = u_2v_j$  where  $j \neq 1, 2$ . Then the label 12 must be assigned to an edge not adjacent to  $e_2$  and  $e_5$ . Let  $f(u_3v_k) = 12$  where  $k \neq 2, j$ .

If  $k = 1$ , since  $e_2$  and  $u_3v_k$  are nonadjacent, the edge  $e_5 = u_2v_j$  with label

10 and the edge  $e_3 = u_1v_3$  with label 4 are adjacent. Thus  $j = 3$ . Now  $w((u_2, v_3, u_1, v_1)) = w((u_3, v_1, u_1, v_2)) = 15$ , a contradiction. If  $k \neq 1$ , then  $w((v_1, u_1, v_2)) + f(e_5) = f(u_1v_1) + f(u_3v_k) = 13$  and hence path weight 13 cannot be obtained. Hence the labels 1 and 2 cannot be assigned to the adjacent edges  $e_1 = u_1v_1$  and  $e_2 = u_1v_2$ .

**Case 2.**  $e_1 = u_1v_1$  and  $e_2 = u_2v_1$ .

Let  $f(e_3) = 4$ . If  $e_3$  is nonadjacent to both  $e_1$  and  $e_2$ , then by Corollary 1, the label 5 must be assigned to an edge nonadjacent to  $e_1$  and  $e_3$ . Thus 1 and 5 are assigned to two nonadjacent edges and 2 and 4 are assigned to two nonadjacent edges. Hence path weight 6 cannot be obtained.

If  $e_3$  is adjacent to  $e_2$  and nonadjacent to  $e_1$ , then 5 must be assigned to an edge nonadjacent to both  $e_1$  and  $e_3$ . Let  $f(u_3v_3) = 5$ . Then the paths  $(u_1, v_1, u_3, v_3)$  and  $(v_2, u_2, v_1, v_3)$  have weight  $6 + f(u_3v_1)$  which is a contradiction.

Now suppose  $e_3$  is adjacent to  $e_1$  and nonadjacent to  $e_2$ . Let  $e_3 = u_1v_2$ . By Corollary 1, the label 6 must be assigned to an edge not adjacent to  $e_2$  and  $e_3$ . Hence  $[7] \subseteq S$ . Again by Corollary 1, the label 8 must be assigned to an edge not adjacent to the edges with labels 2 and 6. But now 2 and 8 are assigned to two nonadjacent edges and 4 and 6 are assigned to nonadjacent edges. Hence by Corollary 3, path weight 10 cannot be obtained.


Therefore  $e_3$  is adjacent to both  $e_1$  and  $e_2$ . Let  $e_3 = u_3v_1$ . Hence  $[6] \subseteq S$ . Let  $f(e_4) = 7$ . If  $e_4$  is nonadjacent to  $e_1$ , then by Corollary 1, the label 8 must be assigned to an edge  $e$  nonadjacent to  $e_1$  and  $e_4$ . Now, if  $e_4$  is nonadjacent to  $e_2$ , then  $f(e_2) + f(e_4) = f(e_1) + f(e) = 9$  and by Corollary 3, the path weight 9 cannot be obtained. If  $e_4$  is adjacent to  $e_2$ , then the label 8 must be assigned to an edge nonadjacent to both  $e_1$  and  $e_4$ . Let  $f(u_3v_3) = 8$ . In this case we get two paths with weight 13. Hence  $e_4$  is adjacent to  $e_1$ . Now if  $e_4$  is nonadjacent to  $e_2$ , then by Corollary 1, the label 9 must be assigned to an edge  $e$  nonadjacent to both  $e_2$  and  $e_4$ . Hence  $f(e_3) + f(e_4) = f(e) + f(e_2) = 11$  and by Corollary 3, the path weight 11 cannot be obtained, which is a contradiction. Hence  $K_{3,n}$  is not a Leech graph.  $\square$

**Corollary 5.** *The Leech index of  $K_{3,n}$  is  $k(K_{3,n}) = 14$ , for  $n \geq 4$ . When  $n = 3$ ,  $k(K_{3,3}) = 10$ .*

### 3. Leech Cycles

In [12], it has been proved that the cycle  $C_n$ , with  $3 \leq n \leq 6$  is a Leech graph. The Leech labelings of these cycles are given in Figure 2, in which two Leech labelings are given for  $C_4$ . Thus for a Leech graph, the Leech labeling is not in general unique. In a cycle, since there exist exactly two paths between every pair of vertices,  $t_p(C_n) =$

$n(n - 1)$ . In this section, we present results on the maximum label that can assigned to an edge in a Leech cycle.



**Figure 2.** Leech labeling of cycles of order  $\leq 6$

**Theorem 3.** Let  $f$  be the Leech labeling of a cycle  $C_n$ . Then,  $w(f) = 2\binom{n}{2} + 1$ .

*Proof.* Let  $C_n = (v_1, v_2, \dots, v_n, v_1)$  and  $f(v_1v_2) = 1$ . Then  $P = (v_2, v_3, \dots, v_n, v_1)$  is the path of maximum weight. Hence  $w(P) = t_p(G) = n(n - 1)$ . Therefore,  $w(f) = w(P) + 1 = n^2 - n + 1 = 2\binom{n}{2} + 1$ .  $\square$

**Theorem 4.** The maximum value that can be assigned to an edge in a Leech labeling of a cycle  $C_n$  is  $\binom{n}{2} + 1$ . Also, this maximum value is attained only by  $C_3$  and  $C_4$ .

*Proof.* Let  $f(e) = M$ , where  $M$  is the maximum value assigned to an edge by  $f$ . Now,  $P_n = C_n - e$  is a path of order  $n$  and hence  $w(C_n - e) \geq 1 + 2 + \dots + (n - 1) = \binom{n}{2}$ . Hence,  $w(f) = 2\binom{n}{2} + 1 = w(C_n - e) + M \geq \binom{n}{2} + M$ . Therefore,  $M \leq \binom{n}{2} + 1$ . Also, equality holds if and only if  $w(P_n) = \binom{n}{2}$  and the path weights of all subpaths of  $P_n$  are exactly  $\{1, 2, \dots, \binom{n}{2} - 1\}$ . Hence  $P_n$  is a Leech path. Since  $P_2, P_3$  and  $P_4$  are the only Leech paths, the maximum edge label  $M$  is attained only for  $C_3$  and  $C_4$ .  $\square$

**Theorem 5.** The only Leech cycles which admits a Leech labeling in which the maximum label is  $\binom{n}{2}$  are  $C_4$  and  $C_5$ .

*Proof.* Let  $f(e_1) = M = \binom{n}{2}$ , where  $M$  is the maximum value assigned to an edge by  $f$  and  $e_1 = v_1v_2$ . Let  $P_n = C_n - e_1$ . Then,  $w(P_n) = \binom{n}{2} + 1$  and all path weights  $1, 2, \dots, \binom{n}{2} - 1$  must be obtained from  $P_n$ . Hence, the set of edge labels of  $P_n$  is  $\{1, 2, \dots, n - 2, n\}$ . Let  $f(e_2) = 1$ . If  $e_2 = v_2v_3$ , then  $w(v_3, v_4, \dots, v_n, v_1) = \binom{n}{2} = w(v_1v_2)$ . Hence,  $e_2 \neq v_2v_3$ . Similarly,  $e_2 \neq v_nv_1$ . If  $e_2 = v_iv_{i+1}$  and  $f(v_{i-1}v_i) = k$  then  $w(v_{i-1}, v_i, v_{i+1}) = k + 1$ . Since  $1, 2, \dots, n - 2$  and  $n$  are already edge weights,  $k \notin \{2, 3, \dots, n - 3\}$ . A similar argument holds for  $f(v_{i+1}v_{i+2}) = l$  also. Therefore,  $k$  and  $l$  are  $n - 2$  and  $n$  in some order. If  $n = 4$ , this gives a Leech labeling of  $C_4$  with maximum label 6 as given in Figure 2.

Now, let  $n \geq 5$ . Let  $f(e_3) = 2$ . If  $e_3$  is adjacent to an edge labeled  $k$  then these two edges together gives a path of weight  $k + 2$ . Since,  $1, 2, \dots, n - 2$  and

$n$  are already edge weights and  $\{w(v_{i-1}v_iv_{i+1}), w(v_iv_{i+1}v_{i+2})\} = \{n-1, n+1\}$ ,  $k \notin \{1, 3, \dots, n-2\}$ . Hence,  $e_3$  is adjacent only to the edge with label  $n$ .

Now, since the edges with labels 1 and 2 are non-adjacent, there exists an edge  $e_4$  with  $f(e_4) = 3$ . If  $e_4$  is adjacent to an edge labeled  $k \in \{1, 2, 4, \dots, n-2\}$  then the path weight of  $e_4$  together with this edge will be in  $\{4, 5, 7, \dots, n+1\}$  which is not possible. Hence, the only possibility is  $n-2 = 3$  and this gives a Leech labeling of  $C_5$  with the maximum edge label is 10 as given in Figure 2.  $\square$

**Theorem 6.** *The only Leech cycle which admits a Leech labeling in which the maximum label is  $\binom{n}{2} - 1$  is  $C_6$ .*

*Proof.* Let  $f(e) = \binom{n}{2} - 1$ , where  $e = v_1v_2$ . Let  $P_n = C_n - e = (v_2, v_3, \dots, v_n, v_1)$ . Then  $w(P_n) = \binom{n}{2} + 2$  and all path weights less than  $\binom{n}{2} - 1$  must be obtained from  $P_n$ . Hence the set of all path weights of the subpaths of  $P_n$  is  $\{1, 2, \dots, \binom{n}{2} - 2, \binom{n}{2} + 2, k\}$ , where  $k$  is  $\binom{n}{2}$  or  $\binom{n}{2} + 1$ . Also, the sum of edge weights of  $P_n$  is  $\binom{n}{2} + 2$  and hence the set of all edge labels of  $P_n$  is  $\{1, 2, \dots, n-2, n+1\}$  or  $\{1, 2, \dots, n-3, n-1, n\}$ . We consider four cases.

**Case 1.**  $k = \binom{n}{2}$  and the set of edge labels of  $P_n$  is  $\{1, 2, \dots, n-2, n+1\}$ . Since  $w(P_n) = \binom{n}{2} + 2$  and  $k = \binom{n}{2}$  is a path weight of a subpath of  $P_n$ , the label 2 must be assigned to a pendant edge of  $P_n$ . Let  $f(v_2v_3) = 2$ . Now since  $f(e) = \binom{n}{2} - 1$  and  $\binom{n}{2}$  is a path weight of a subpath of  $P_n$ , the label 1 cannot be assigned to a pendant edge of  $P_n$ . Let  $f(e_1) = 1$ , where  $e_1$  is an internal edge of  $P_n$ . Now, if  $a$  and  $b$  are edge labels of two adjacent edges of  $P_n$ , then  $a + b$  is not an edge label. Hence,  $a + b = n - 1$  or  $n$  or  $a + b \geq n + 2$ . Hence the two edges adjacent to  $e_1$  have labels  $n - 2$  and  $n + 1$  and we get path weights  $n - 1$  and  $n + 2$ . Now, if  $f(v_3v_4) = x$  then,  $w(v_2, v_3, v_4) = x + 2 = n$  or  $x + 2 \geq n + 3$ . If  $x + 2 = n$ , then  $x = n - 2$  and hence  $e_1 = v_4v_5$ . But, then  $w(v_2, v_3, v_4, v_5) = 2 + n - 2 + 1 = n + 1$  which is already an edge weight, a contradiction. Therefore, the only possibility is  $f(v_3v_4) = n + 1$  and eventually  $f(v_4v_5) = 1$  and  $f(v_5v_6) = n - 2$  and we get path weights  $n + 3$  and  $n + 4$ . Now, there is an edge with label 3 and if the edge adjacent to it is labeled  $y$  then  $y + 3 = n$  or  $y + 3 \geq n + 5$ . But,  $y \geq n + 2$  is not possible and hence 3 is assigned to a pendant edge of  $P_n$ . But, then the path weight  $3 + \binom{n}{2} - 1 = \binom{n}{2} + 2$  repeats. Hence, this case is not possible.

**Case 2.**  $k = \binom{n}{2}$  and the set of edge labels of  $P_n$  is  $\{1, 2, \dots, n-3, n-1, n\}$ . As in Case 1,  $f(v_2v_3) = 2$  and  $f(e_1) = 1$ , where  $e_1$  is an internal edge of  $P_n$ . Also, if  $a$  and  $b$  are edge labels of two adjacent edges of  $P_n$ , then  $a + b = n - 2$  or  $a + b \geq n + 1$ . Hence the two edges adjacent to  $e_1$  have labels  $n - 3$  and  $n$  and we get path weights  $n - 2$  and  $n + 1$ . Now if  $f(v_3v_4) = x$  then,  $w(v_2, v_3, v_4) = x + 2 \geq n + 2$ . Therefore,  $f(v_3v_4) = n$  and eventually,  $f(v_4v_5) = 1$  and  $f(v_5v_6) = n - 3$ . Also, we have obtained all path weights up to  $n + 3$ . Now, if the label  $y$  is assigned to an edge adjacent to the edge labeled 3, then  $y + 3 \geq n + 4$  which is not possible.




**Case 3.**  $k = \binom{n}{2} + 1$  and the set of edge labels of  $P_n$  is  $\{1, 2, \dots, n-2, n+1\}$ . Since  $w(P_n) = \binom{n}{2} + 2$  and  $k = \binom{n}{2} + 1$  is a path weight of a subpath of  $P_n$ , the label 1 must be assigned to a pendant edge of  $P_n$ . Let  $f(v_2v_3) = 1$ . Also, since  $\binom{n}{2}$  is not a path weight of a subpath of  $P_n$ , 2 cannot be assigned to a pendent edge of  $P_n$ . Let  $f(e_1) = 2$ , where  $e_1$  is an internal edge of  $P_n$ . Now, if  $a$  and  $b$  are edge labels of two adjacent edges of  $P_n$ , then  $a + b = n - 1$  or  $n$  or  $a + b \geq n + 2$ . Therefore,  $f(v_3v_4) = n - 2$  or  $n + 1$ . If  $f(v_3v_4) = n - 2$  then we get path weight  $n - 1$  also, so that if an edge adjacent to  $e_1$  is given label  $x$ , then  $x + 2 = n$  or  $x + 2 \geq n + 2$ . Therefore, the edges adjacent to  $e_1$  are labeled  $n - 2$  and  $n + 1$ . Therefore,  $f(v_4v_5) = 2$  and  $f(v_5v_6) = n + 1$ . But, then  $w(v_2, v_3, v_4, v_5) = 1 + n - 2 + 2 = n + 1$ , which is already an edge weight. Therefore, let  $f(v_3v_4) = n + 1$  and then we get the path weight  $n + 2$  also. Again, if an edge adjacent to  $e_1$  is given label  $x$ , then  $x + 2 = n - 1$  or  $n$  or  $x + 2 \geq n + 3$ . Therefore, the edges adjacent to  $e_1$  are labeled  $n - 3$  or  $n - 2$  or  $n + 1$ . If  $e_1$  is adjacent to an edge labeled  $n + 1$  then,  $e_1 = v_4v_5$  and we get paths of weight  $n + 3$  and  $n + 4$ . Now, the edge labeled 3 can be adjacent only to either  $n - 3$  or the edge labeled  $n - 4$ , which implies 3 is assigned to a pendant edge of  $P_n$ . But, then  $f(e) + 3$  gives another path weight of weight  $\binom{n}{2} + 2$ , a contradiction. Therefore, the labels of edges adjacent to 2 are  $n - 3$  and  $n - 2$ , so that the path weights  $n - 1$  and  $n$  are obtained. Again, if the label  $y$  is assigned to an edge adjacent to the edge labeled 3, then  $y + 3 \geq n + 3$  in which case also 3 is assigned to a pendant edge and hence is not possible.

**Case 4.**  $k = \binom{n}{2} + 1$  and the set of edge labels of  $P_n$  is  $\{1, 2, \dots, n - 3, n - 1, n\}$ . As in Case 3,  $f(v_2v_3) = 1$  and  $f(e_1) = 2$ , where  $e_1$  is an internal edge of  $P_n$ . Again, if  $a$  and  $b$  are edge labels of two adjacent edges of  $P_n$ , then  $a + b = n - 2$  or  $a + b \geq n + 1$ . Therefore,  $f(v_3v_4) = n - 3$  or  $n$ .

If  $f(v_3v_4) = n$ , then we get path weight  $n + 1$  also. Now, if an edge adjacent to  $e_1$  is given label  $x$ , then  $x + 2 = n - 2$  or  $x + 2 \geq n + 2$ . Therefore, the edges adjacent to 2 are labeled  $n - 4$  and  $n$ , so that we get all path weights upto  $n + 3$ . Now, if label  $y$  is assigned to an edge adjacent to the edge labeled 3, then  $y + 3 \geq n + 4$  which is not possible. Therefore, let  $f(v_3v_4) = n - 3$  so that we get path weight  $n - 2$  also, so that if an edge adjacent to  $e_1$  is given label  $x$ , then  $x + 2 \geq n + 1$ . Therefore, the edges adjacent to  $e_1$  are labeled  $n - 1$  and  $n$ , so that we get all path weights up to  $n + 2$ . In this case also, if the label  $y$  is assigned to an edge adjacent to the edge labeled 3, then  $y + 3 \geq n + 3$  in which case 3 is assigned to a pendant edge and hence is not possible. So, the only possibility is  $n - 3 = 3$  and this gives the Leech labeling of  $C_6$ .  $\square$

**Corollary 6.** *The maximum value that can be assigned to an edge in a Leech labeling of a cycle  $C_n$ , where  $n > 7$ , is less than  $\binom{n}{2} - 1$ .*

The following figure gives three different Leech labelings of  $C_6$  in which the maximum labels are  $\binom{n}{2} - 1$ ,  $\binom{n}{2} - 2$  and  $\binom{n}{2} - 3$ . We strongly believe that cycles of length greater than 6 are not Leech graphs.



**Figure 3.** Leech labelings of  $C_6$

**Acknowledgement.** The second author thank Cochin University of Science and Technology for granting project under Seed Money for New Research Initiatives (order No.CUSAT/PL(UGC).A1/1112/2021) dated 09.03.2021. The authors are thankful to an anonymous reviewer, whose helpful suggestions resulted is substantial improvement in the paper.

**Conflict of interest.** The authors declare that they have no conflict of interest.

**Data Availability.** Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

## References

- [1] B. Calhoun, K. Ferland, L. Lister, and J. Polhill, *Minimal distinct distance trees*, J. Combin. Math. Combin. Comput. **61** (2007), 33–57.
- [2] G. Chartrand, L. Lesniak, and P. Zhang, *Graphs & Digraphs*, vol. 22, Chapman & Hall London, 1996.
- [3] D. Leach, *Modular Leech trees of order at most 8*, Int. J. Combin. **2014** (2014), Article ID: 218086.  
<https://doi.org/10.1155/2014/218086>.
- [4] D. Leach and M. Walsh, *Generalized leech trees*, J. Combin. Math. Combin. Comput. **78** (2011), 15–22.
- [5] J. Leech, *Another tree labeling problem*, Amer. Math. Monthly **82** (1975), no. 9, 923–925.
- [6] M. Ozen, H. Wang, and D. Yalman, *Note on Leech-type questions of tree*, Integers **16** (2016), #A21.
- [7] L.A. Szekely, H. Wang, and Y. Zhang, *Some non-existence results on Leech trees*, Bull. Inst. Combin. Appl. **44** (2005), 37–45.
- [8] H. Taylor, *Odd path sums in an edge-labeled tree*, Math. Magazine **50** (1977), no. 5, 258–259.  
<https://doi.org/10.1080/0025570X.1977.11976658>.
- [9] ———, *A distinct distance set of 9 nodes in a tree of diameter 36*, Discrete Math. **93** (1991), no. 2-3, 167–168.  
[https://doi.org/10.1016/0012-365X\(91\)90252-W](https://doi.org/10.1016/0012-365X(91)90252-W).

- 
- [10] S. Varghese, A. Lakshmanan S., and S. Arumugam, *Two classes of non-Leech trees*, Electron. J. Graph Theory Appl. **8** (2020), no. 1, 205–210.  
<https://doi.org/10.5614/ejgta.2020.8.1.15>.
- [11] ———, *Leech index of a tree*, J. Discrete Math. Sci. Crypt. **25** (2022), no. 8, 2237–2247.  
<https://doi.org/10.1080/09720529.2020.1800217>.
- [12] ———, *Two extensions of Leech labeling to the class of all graphs*, AKCE Int. J. Graphs Comb. **19** (2022), no. 2, 159–165.  
<http://doi.org/10.1080/09728600.2022.2084354>.
- [13] ———, *Leech labeling problem on tristar*, AIP Conf. Proc. **2649** (2023), no. 1, Article ID: 020007.  
<https://doi.org/10.1063/5.0114834>.