

Edge metric dimension of silicate networks

Savari Prabhu[†], T. Jenifer Janany^{*}

Department of Mathematics, Rajalakshmi Engineering College, Chennai 602105, India
[†]drsavariprabhu@gmail.com
^{*}jeniferjanany@gmail.com

Received: 18 March 2024; Accepted: 7 July 2024
Published Online: 20 July 2024

Abstract: Metric dimension is an essential parameter in graph theory that aids in addressing issues pertaining to information retrieval, localization, network design, and chemistry through the identification of the least possible number of elements necessary to identify the vertices in a graph uniquely. A variant of metric dimension, called the edge metric dimension focuses on distinguishing the edges in a graph G , with a vertex subset. The minimum possible number of vertices in such a set is denoted as $\dim_E(G)$. This paper presents the precise edge metric dimension of silicate networks.

Keywords: edge metric basis, silicate, twins, tetrahedron, chain silicate, cyclic silicate.

AMS Subject classification: 05C69, 05C12

1. Introduction

Metric dimension is a measure of how efficiently one can locate and distinguish between the vertices (nodes) of a graph using a minimal set of landmarks or reference points [5, 23]. Metric dimension has applications in diverse domains, particularly network design [3], chemistry [4, 7], robotics [9], and location-based services. Calculating the exact metric dimension of a graph is often a computationally challenging problem. For particular class of graphs, such as trees, there are efficient algorithms to find the metric dimension. However, for general graphs [9], bipartite graphs [16] and directed graphs [22], this problem remains NP-hard. Despite the computational difficulty, the precise value of metric dimension is evaluated for many graph structures

^{*} *Corresponding Author*

including honeycomb [15], TiO_2 nanotubes [19], butterfly [16], benes [16], Sierpiński [10], silicate [17], and irregular triangular networks [21].

The distance $d_G(u, v)$ between two vertices u and v in a connected graph G , where $V(G)$ is the set of vertices and $E(G)$ is the set of edges, is defined as the minimum number of edges in any shortest path (geodesic) connecting them. The distance between an edge $e = vw$ and a vertex u is given by, $d_G(e, u) = \min\{d_G(v, u), d_G(w, u)\}$. Based on the distance from vertices of an ordered subset $X = \{u_1, u_2, \dots, u_l\}$, every vertex v is represented with a vector of distance,

$$r(v|X) = (d_G(v, u_1), d_G(v, u_2), \dots, d_G(v, u_l)).$$

The subset X is a resolving set (**RS**) if $r(x|X) \neq r(y|X)$, $\forall x, y \in V(G)$. To put it in another way, X is a **RS** if for every $x, y \in V(G)$, a vertex $u \in X$ exists such that distances from x and y to u are unequal. See Figure 1(a). We use the term (metric) basis to represent a **RS** with minimum vertices and metric dimension (**MD**), $\dim(G)$ to indicate its cardinality.




Figure 1. (a) Resolving set (**RS**); (b) Edge resolving set (**ERS**)

The vertices of a basis distinguish the vertices in the graph and not necessarily the edges. This fact ignited the spark of edge metric basis (**EMB**). A vertex subset S , is called an edge resolving set (**ERS**) when we have a vertex $w \in S$ for every pair of edges e, f , such that $d_G(w, e)$ and $d_G(w, f)$ are different. See Figure 1(b). **EMB** is a vertex subset which resolves the edges of the graph with minimum possible vertices and the cardinality of **EMB** is the edge metric dimension (**EMD**), $\dim_E(G)$. This idea was conceptualized in [8] by Kelenc et al. and is concluded as NP-complete. In the aftermath of this concept's origin a number of articles has outbroken in this field. To list a few, we have the characterization of graphs with maximum **EMD** [27, 28], the **EMD** of grid graphs [8], web graph, prism related graph, convex polytope antiprism, convex polytope graph [26], benzenoid tripod structure [2], honeycomb network, hexagonal network [1], Erdos-Renyi random graph [29], and generalized Petersen graph $P(n, 3)$ [25]. The research regarding this realm has expanded to graph operations such as join, lexicographic, corona [18], and hierarchical products [11]. Identifying graphs with **EMD** lesser than **MD** received specific attention [12, 13].

2. Silicate Networks

Silicates are among the most abundant minerals on Earth and are the primary components of many materials, including rocks, minerals, and glasses. A silicon atom joined with equally spaced oxygen atoms when oxygen atoms are placed at the four corners of a tetrahedron produces a silicate tetrahedron. The basic building block of silicate networks is the silicate tetrahedron, SiO_4 . See Figure 2. In a silicate network (or sheet), each oxygen atom is shared between adjacent tetrahedra, leading to the formation of a continuous network. The silicate network was perceived as an interconnection network, inspired by the molecular structure of SiO_4 . Various properties, including metric dimension [17], topological descriptors [6, 14], power domination [24], and fault-tolerant metric dimension [20] have been discussed.




Figure 2. Silicate tetrahedron

When a silicate tetrahedron connects with other tetrahedra linearly, a single-row silicate chain is created. A chain silicate with n tetrahedra is denoted by CS_n . It has $3n + 1$ vertices and $6n$ edges. See figure 3.




Figure 3. Chain silicate CS_7

Cyclic silicates are a specific type of silicate mineral or compound characterized by a ring-like structure. In cyclic silicates, the silicate tetrahedra are arranged in closed loops or rings, giving rise to distinct structural and chemical properties. A cyclic silicate CC_n can be obtained from a cycle of length n , by replacing each edge with a tetrahedron. CC_n has $3n$ vertices and $6n$ edges. See Figure 4.




Figure 4. Cyclic silicate CC_8

3. Main Results

A one point union of two tetrahedrons is called a twin tetrahedron in silicate network. In CS_n we have twin tetrahedrons with five 3-degree vertices and some with four 3-degree vertices. See Figure 5.




Figure 5. Twin tetrahedrons

Let us define a cubic set, C to denote the set of 3-degree vertices in a twin tetrahedron. We have the following lemma regarding the role of cubic vertices in **ERS** of silicate networks.

Lemma 1. *Let G be any silicate network and S be an **ERS** of G . Then for every twin tetrahedron at most one vertex of C can be excluded from S i.e., $|C \setminus S| \leq 1$.*

Proof. Suppose there exists a twin tetrahedron with two cubic vertices $p, q \notin S$. Let v_0 be the intersection point of the twin tetrahedron and let $e_1 = v_0p$, $e_2 = v_0q$. Then $d(e_1, u) = d(e_2, u) = d(v_0, u)$ for each $u \in V(G) \setminus \{p, q\}$, which contradicts S being an **ERS**. \square

Similar to the twin tetrahedrons, we have two types of tetrahedrons, some with three cubic vertices (Type I) and the remaining with two cubic vertices (Type II). See Figure 6.




Figure 6. Tetrahedrons in silicate network Type I and Type II

Consider a type I tetrahedron. Let v_0, v_1, v_2, v_3 be the vertices and the edges of the tetrahedron be $e_1 = v_0v_1, e_2 = v_0v_2, e_3 = v_0v_3, e_4 = v_1v_2, e_5 = v_1v_3, e_6 = v_2v_3$. Let $S_1 = \{s, v_1, v_2\}$ and $d(s, v_1) = x$. Then

$$\begin{aligned} r(e_1|S_1) &= (x, 0, 1) \\ r(e_2|S_1) &= (x + 1, 1, 0) \\ r(e_3|S_1) &= (x + 1, 1, 1) \\ r(e_4|S_1) &= (x, 0, 0) \\ r(e_5|S_1) &= (x, 0, 1) \\ r(e_6|S_1) &= (x + 1, 1, 0) \end{aligned}$$

Similarly, in type II tetrahedron let u_0, u_1, u_2, u_3 be the vertices and $e_1 = u_0u_1, e_2 = u_0u_2, e_3 = u_0u_3, e_4 = u_1u_2, e_5 = u_1u_3, e_6 = u_2u_3$ be the edges. If $S_2 = \{s, t, u_2\}$, $d(s, u_0) = x$ and $d(t, u_1) = y$ then

$$\begin{aligned} r(e_1|S_2) &= (x, y, 1) \\ r(e_2|S_2) &= (x + 1, y + 1, 0) \\ r(e_3|S_2) &= (x + 1, y + 1, 1) \\ r(e_4|S_2) &= (x, y, 0) \\ r(e_5|S_2) &= (x, y, 1) \\ r(e_6|S_2) &= (x + 1, y + 1, 0) \end{aligned}$$

The following remark and lemma are straightforward from the above observations.

Remark 1. For tetrahedrons of type I, $|S \cap C| \geq 2$ and for type II, $|S \cap C| \geq 1$, where S is an **ERS** and C is the set of 3-degree vertices.

Lemma 2. In a silicate network G , if for a vertex subset S , the following conditions holds, then S is an **ERS**.

- (i) For every twin tetrahedron $|C \setminus S| \leq 1$
- (ii) For every type I tetrahedron, $|S \cap C| \geq 2$

(iii) For every type II tetrahedron, $|S \cap C| \geq 1$

Theorem 1. For any even integer $n \geq 2$, $\dim_E(CS_n) = \frac{3n}{2} + 2$.

Proof. When n is even, CS_n can be partitioned into $\frac{n}{2}$ edge-disjoint twin tetrahedrons. Let $C_1, C_2, \dots, C_{\frac{n}{2}}$ be the corresponding cubic sets. Since $|C_1| = |C_{\frac{n}{2}}| = 5$ and $|C_2| = \dots = |C_{\frac{n}{2}-1}| = 4$, by Lemma 1, $\dim_E \geq 3(\frac{n}{2} - 2) + 4(2) = \frac{3n}{2} + 2$.

Consider CS_n as a path on n vertices $\{w_1, w_2, \dots, w_n\}$. Define a mapping $l : \{w_1, w_2, \dots, w_n\} \rightarrow \{1, 2\}$ as follows,

$$l(w_1) = l(w_2) = l(w_{n-1}) = l(w_n) = 2$$

for $3 \leq i \leq n - 2$,

$$l(w_i) = \begin{cases} 1, & i \equiv 1 \pmod{2} \\ 2, & i \equiv 0 \pmod{2}. \end{cases}$$

Consider a collection of 3-degree vertices S , with the property that $|S \cap R_i| = l(w_i)$, where R_i is the set of vertices of the tetrahedron corresponding to w_i . By Lemma 2, S is an **ERS** of CS_n and $|S| = \frac{3n}{2} + 2$. Thus, $\dim_E(CS_n) = \frac{3n}{2} + 2$. \square

Theorem 2. For any odd integer $n \geq 1$, $\dim_E(CS_n) = \frac{3(n+1)}{2}$.

Proof. CS_n with n odd, has two twin tetrahedrons with five cubic vertices, $\frac{n-1}{2} - 2$ twin tetrahedrons with four cubic vertices and a single tetrahedron with two cubic vertices. By Lemma 1, $\dim_E(CS_n) \geq \frac{3(n+1)}{2}$.

Define a mapping $l : \{w_1, w_2, \dots, w_n\} \rightarrow \{1, 2\}$ as follows,

$$l(w_1) = l(w_2) = l(w_{n-1}) = l(w_n) = 2$$

and for $3 \leq i \leq n - 2$,

$$l(w_i) = \begin{cases} 1, & i \equiv 1 \pmod{2} \\ 2, & i \equiv 0 \pmod{2}. \end{cases}$$

Here each w_i represents a tetrahedron of CS_n . Let S be a set of 3-degree vertices with $|S \cap R_i| = l(w_i)$, where R_i is the set of vertices of the tetrahedron corresponding to w_i . Since S satisfies the conditions in Lemma 2, and $|S| = \frac{3(n+1)}{2}$, it follows that $\dim_E = \frac{3(n+1)}{2}$. \square

Theorem 3. For any even integer $n \geq 4$, $\dim_E(CC_n) = \frac{3n}{2}$.

Proof. For even values of n , CC_n has $\frac{n}{2}$ twin tetrahedrons that are edge-disjoint and each cubic set has four vertices. Thus by Lemma 1, $\dim_E \geq \frac{3n}{2}$. Consider CC_n as a cycle on n vertices $\{w_1, w_2, \dots, w_n\}$. Define a mapping $l : \{w_1, w_2, \dots, w_n\} \rightarrow \{1, 2\}$ as follows, for $1 \leq i \leq n$,

$$l(w_i) = \begin{cases} 1, & i \equiv 1 \pmod{2} \\ 2, & i \equiv 0 \pmod{2}. \end{cases}$$

Let S be a set of 3-degree vertices with $|S \cap R_i| = l(w_i)$, where R_i is the set of vertices of the tetrahedron corresponding to w_i . By Lemma 2, S resolves the edges of CC_n . Here $|S| = \frac{3n}{2}$ and this implies that, $\dim_E(CC_n) = \frac{3n}{2}$. \square

Theorem 4. For any odd integer $n \geq 3$, $\dim_E(CC_n) = \frac{3(n+1)}{2} - 1$.

Proof. In CC_n , when n is odd, we have $\frac{n-1}{2}$ twin tetrahedron with four cubic vertices and a single tetrahedron. Thus, $\dim_E(CC_n) \geq \frac{3(n+1)}{2} - 1$. Consider CC_n as a cycle on n vertices $\{w_1, w_2, \dots, w_n\}$. Define a mapping $l : \{w_1, w_2, \dots, w_n\} \rightarrow \{1, 2\}$ as follows, for $1 \leq i \leq n-1$,

$$l(w_i) = \begin{cases} 1, & i \equiv 1 \pmod{2} \\ 2, & i \equiv 0 \pmod{2}, \end{cases}$$

and $l(w_n) = 2$. Consider a collection of 3-degree vertices S , with the property that $|S \cap R_i| = l(w_i)$, where R_i is the set of vertices of the tetrahedron corresponding to w_i . By Lemma 2, S is an edge resolving set of CC_n and since $|S| = \frac{3(n+1)}{2} - 1$, we get $\dim_E(CC_n) = \frac{3(n+1)}{2} - 1$. \square

4. Conclusion

Silicate networks are prevalent in nature, forming the backbone of many minerals in the Earth's crust. They also have significant industrial applications, particularly in the production of glasses, ceramics, and construction materials. Understanding the structure and properties of silicate networks is essential for designing and engineering materials with specific functionalities and characteristics. This study has identified the necessary vertices to resolve the edges in cyclic and chain silicate networks. Future works can focus on this parameter for other types of silicate networks.

Acknowledgements: The authors are thankful to Rajalakshmi Educational Trust for granting financial support under Centre for Sponsored Research and Consultancy/CSRC84.

Conflict of Interest: The authors declare that they have no conflict of interest.

Data Availability: Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

References

- [1] S. Abbas, Z. Raza, N. Siddiqui, F. Khan, and T. Whangbo, *Edge metric dimension of honeycomb and hexagonal networks for IoT*, *Comput. Mater. Contin.* **71** (2022), no. 2, 2683–2695.
- [2] A. Ahmad, S. Husain, M. Azeem, K. Elahi, and M.K. Siddiqui, *Computation of edge resolvability of benzenoid tripod structure*, *J. Math.* **2021** (2021), no. 1, Article ID: 9336540.
<https://doi.org/10.1155/2021/9336540>.
- [3] Z. Beerliova, F. Eberhard, T. Erlebach, A. Hall, M. Hoffmann, M. Mihal'ak, and L.S. Ram, *Network discovery and verification*, *IEEE J. Sel. Areas Commun.* **24** (2006), no. 12, 2168–2181.
<https://doi.org/10.1109/JSAC.2006.884015>.
- [4] G. Chartrand, L. Eroh, M.A. Johnson, and O.R. Oellermann, *Resolvability in graphs and the metric dimension of a graph*, *Discrete Appl. Math.* **105** (2000), no. 1-3, 99–113.
[https://doi.org/10.1016/S0166-218X\(00\)00198-0](https://doi.org/10.1016/S0166-218X(00)00198-0).
- [5] F. Harary and R.A. Melter, *On the metric dimension of a graph*, *Ars Combin.* **2** (1976), 191–195.
- [6] S. Hayat and M. Imran, *Computation of topological indices of certain networks*, *Appl. Math. Comput.* **240** (2014), 213–228.
<https://doi.org/10.1016/j.amc.2014.04.091>.
- [7] M. Johnson, *Structure-activity maps for visualizing the graph variables arising in drug design*, *J. Biopharm. Stat.* **3** (1993), no. 2, 203–236.
<https://doi.org/10.1080/10543409308835060>.
- [8] A. Kelenc, N. Tratnik, and I.G. Yero, *Uniquely identifying the edges of a graph: The edge metric dimension*, *Discrete Appl. Math.* **251** (2018), 204–220.
<https://doi.org/10.1016/j.dam.2018.05.052>.
- [9] S. Khuller, B. Raghavachari, and A. Rosenfeld, *Landmarks in graphs*, *Discrete Appl. Math.* **70** (1996), no. 3, 217–229.
[https://doi.org/10.1016/0166-218X\(95\)00106-2](https://doi.org/10.1016/0166-218X(95)00106-2).
- [10] S. Klavžar and S.S. Zemljič, *On distances in Sierpiński graphs: Almost-extreme vertices and metric dimension*, *Appl. Anal. Discrete Math.* **7** (2013), no. 1, 72–82.
<https://doi.org/10.2298/AADM130109001K>.
- [11] S. Klavžar and M. Tavakoli, *Edge metric dimensions via hierarchical product and integer linear programming*, *Optim. Lett.* **15** (2021), 1993–2003.

- <https://doi.org/10.1007/s11590-020-01669-x>.
- [12] M. Knor, S. Majstorović, A.T.M. Toshi, R. Škrekovski, and I.G. Yero, *Graphs with the edge metric dimension smaller than the metric dimension*, Appl. Math. Comput. **401** (2021), Article ID: 126076.
<https://doi.org/10.1016/j.amc.2021.126076>.
- [13] M. Knor, R. Škrekovski, and I.G. Yero, *A note on the metric and edge metric dimensions of 2-connected graphs*, Discrete Appl. Math. **319** (2022), 454–460.
<https://doi.org/10.1016/j.dam.2021.02.020>.
- [14] J.B. Liu, S. Wang, C. Wang, and S. Hayat, *Further results on computation of topological indices of certain networks*, IET Control Theory Appl. **11** (2017), no. 13, 2065–2071.
<https://doi.org/10.1049/iet-cta.2016.1237>.
- [15] P. Manuel, R. Bharati, and I. Rajasingh, *On minimum metric dimension of honeycomb networks*, J. Discrete Algorithms **6** (2008), no. 1, 20–27.
<https://doi.org/10.1016/j.jda.2006.09.002>.
- [16] P.D. Manuel, M.I. Abd-El-Barr, I. Rajasingh, and B. Rajan, *An efficient representation of benes networks and its applications*, J. Discrete Algorithms **6** (2008), no. 1, 11–19.
<https://doi.org/10.1016/j.jda.2006.08.003>.
- [17] P.D. Manuel and I. Rajasingh, *Minimum metric dimension of silicate networks*, Ars Combin. **98** (2011), 501–510.
- [18] I. Peterin and I.G. Yero, *Edge metric dimension of some graph operations*, Bull. Malays. Math. Sci. Soc. **43** (2020), no. 3, 2465–2477.
<https://doi.org/10.1007/s40840-019-00816-7>.
- [19] S. Prabhu, T. Flora, and M. Arulperumjothi, *On independent resolving number of $TiO_2[m, n]$ nanotubes*, J. Intell. Fuzzy Syst. **35** (2018), no. 6, 6421–6425.
<https://doi.org/10.3233/JIFS-181314>.
- [20] S. Prabhu, V. Manimozhi, M. Arulperumjothi, and S. Klavžar, *Twin vertices in fault-tolerant metric sets and fault-tolerant metric dimension of multistage interconnection networks*, Appl. Math. Comput. **420** (2022), Article ID: 126897.
<https://doi.org/10.1016/j.amc.2021.126897>.
- [21] S. Prabhu, D. Sagaya Rani Jeba, M. Arulperumjothi, and S. Klavžar, *Metric dimension of irregular convex triangular networks*, AKCE Int. J. Graphs Comb. (2023), In press.
<https://doi.org/10.1080/09728600.2023.2280799>.
- [22] B. Rajan, I. Rajasingh, J.A. Cynthia, and P. Manuel, *Metric dimension of directed graphs*, Int. J. Comput. Math. **91** (2014), no. 7, 1397–1406.
<https://doi.org/10.1080/00207160.2013.844335>.
- [23] P.J. Slater, *Leaves of trees*, Congr. Numer. **14** (1975), 549–559.
- [24] S. Stephen, B. Rajan, J. Ryan, C. Grigorious, and A. William, *Power domination in certain chemical structures*, J. Discrete Algorithms **33** (2015), 10–18.
<https://doi.org/10.1016/j.jda.2014.12.003>.
- [25] D.G.L. Wang, M.M.Y. Wang, and S. Zhang, *Determining the edge metric dimension of the generalized Petersen graph $P(n, 3)$* , J. Comb. Optim. **43** (2022), no. 2,

- 460–496.
<https://doi.org/10.1007/s10878-021-00780-8>.
- [26] Y. Zhang and S. Gao, *On the edge metric dimension of convex polytopes and its related graphs*, J. Comb. Optim. **39** (2020), no. 2, 334–350.
<https://doi.org/10.1007/s10878-019-00472-4>.
- [27] E. Zhu, A. Taranenko, Z. Shao, and J. Xu, *On graphs with the maximum edge metric dimension*, Discrete Appl. Math. **257** (2019), 317–324.
<https://doi.org/10.1016/j.dam.2018.08.031>.
- [28] N. Zubrilina, *On the edge dimension of a graph*, Discrete Math. **341** (2018), no. 7, 2083–2088.
<https://doi.org/10.1016/j.disc.2018.04.010>.
- [29] ———, *Asymptotic behavior of the edge metric dimension of the random graph*, Discuss. Math. Graph Theory **41** (2021), no. 2, 589–599.
<https://doi.org/10.7151/dmgt.2210>.