

## Extremal trees for the general Sombor index

Chinglensana Phanjoubam<sup>1,†</sup>, Sainkupar Mn Mawiong<sup>2,\*</sup>,  
Ardeline M. Buhphang<sup>1,‡</sup>

<sup>1</sup>Department of Mathematics, North-Eastern Hill University, Shillong, India

<sup>†</sup>phanjoubam17@gmail

<sup>‡</sup>ardeline17@gmail

<sup>2</sup>Department of Basic Sciences and Social Sciences, North-Eastern Hill University, Shillong, India

\*skupar@gmail

*Received: 11 February 2024; Accepted: 11 November 2024*

*Published Online: 28 November 2024*

**Abstract:** Recently, the Sombor index of a graph has been extended to general Sombor index. The general Sombor index of a simple graph  $G$  is defined as  $SO_\alpha(G) =$

$\sum_{uv \in E(G)} [d_G(u)^2 + d_G(v)^2]^{\alpha/2}$ , where  $d_G(u)$  denotes the degree of a vertex  $u$  in  $G$  and

$\alpha$  is a real number. In this paper, we obtain bounds for the general Sombor index of trees. We further determine the trees with the extremal general Sombor indices.

**Keywords:** Sombor index, general Sombor index, trees, degree.

**AMS Subject classification:** 05C07, 05C09

### 1. Introduction

One generally associates various graph-theoretical invariants to molecular graphs in chemical graph theory. This way, one studies the correlation of the invariants with various properties of the corresponding molecules. In 1947, Wiener [20] introduced such index for the first time and used it to correlate boiling points of alkanes. Historically, the first vertex-degree-based indices are the Zagreb indices. In general, the vertex-degree-based topological indices of a simple graph  $G$  are of the form

$\sum_{uv \in E(G)} f(d_G(u), d_G(v))$ , where  $d_G(u)$  denotes degree of the vertex  $u$ ,  $E(G)$  is the

edge set and  $f$  is an appropriately chosen function such that  $f(x, y) = f(y, x)$ . A

---

\* Corresponding Author

number of invariants and their generalisations have been introduced in the literature with varying success in quantitative structure-activity relationship (QSAR) and quantitative structure-property relationship (QSPR) studies. In 1975, Randić introduced an index, now known as *Randić connectivity index*, which turned out to be the most widely used index in the QSPR and the QSAR studies [16]. Erdős and Bollobás extended the Randić connectivity index in 1998 [1], known as the *general Randić connectivity index*, and is defined as

$$R_\alpha(G) = \sum_{uv \in E(G)} [d_G(u)d_G(v)]^\alpha$$

for any real number  $\alpha$ . This extends the classical Randić index  $R_{1/2}(G)$  and the second Zagreb index  $R_1(G)$ , another important index that is studied extensively [13]. Extremal results on trees for the general Randić connectivity index are studied extensively in the literature, namely in [3], [9], [8], [10], [12], [21] and [22]. Motivated by Randić connectivity and Zagreb indices, Zhou and Trinajstić defined *sum-connectivity index*  $\chi(G)$  [24] and its generalisation known as *general sum-connectivity index*  $\chi_\alpha(G)$  [25], which are defined as

$$\chi(G) = \sum_{uv \in E(G)} \frac{1}{\sqrt{d_G(u) + d_G(v)}}$$

and

$$\chi_\alpha(G) = \sum_{uv \in E(G)} [d_G(u) + d_G(v)]^\alpha$$

for any real number  $\alpha$ . Notice that  $\chi_{-1/2}(G) = \chi(G)$  and  $\chi_1(G) = M_1(G)$ , the classical first Zagreb index. Thus the general sum-connectivity index extends the ordinary sum-connectivity index and the first Zagreb index. Extremal results on trees for the general sum-connectivity index are explored in [4], [6], [18] and [23].

In this paper, we explore extremal trees for the extension of the recently introduced index, the *Sombor index*, by Gutman [7]. The Sombor index is known to have a reasonable predictive potential in the QSPR and the QSAR studies [17] and its relation with the existing well known indices are studied in [2], [5], [14] and [19]. For extremal results and various bounds of the Sombor index, we refer the readers to a recent survey paper [11] and the references therein.

Recently in [15], the Sombor index is extended, known as the *general Sombor index*, and is defined as

$$SO_\alpha(G) = \sum_{uv \in E(G)} [d_G(u)^2 + d_G(v)^2]^{\alpha/2}$$

for any real number  $\alpha$ . Notice that  $SO_1(G) = SO(G)$ , the usual Sombor index, and  $SO_2(G) = F(G)$ , the forgotten index. Thus the general Sombor index generalises both the ordinary Sombor index and the forgotten index. In [15], bounds in terms

of important graph parameters for general Sombor index have been reported and explored the Nordhaus-Gaddum-type results. It further explored its relations with general sum-connectivity and general Randić indices. In this paper, we derive bounds for the general Sombor index of trees. We further determine the trees with the extremal general Sombor indices.

## 2. Preliminaries

All graphs considered in this paper are finite and simple. For a graph  $G$ , we denote the vertex set of  $G$  by  $V(G)$ . A tree  $T$  is a graph which is connected and acyclic. We denote the  $n$ -vertex path and  $n$ -vertex star by  $P_n$  and  $S_n$  respectively. We now present the following lemmas that are crucial in proving our main results in the next section.

**Lemma 1.** *Let  $H$  be a connected graph of order at least 2. Let  $a_1 \geq a_2 \geq 1$  and let  $v \in V(H)$ . Let  $H_1$  be the graph obtained from  $H$  by attaching two paths  $P$  and  $P'$  of lengths  $a_1$  and  $a_2$  respectively to  $v$ ; and  $H_2$  be the graph obtained from  $H$  by attaching a path  $P''$  of length  $a_1 + a_2$  to  $v$ . Then  $SO_\alpha(H_1) > SO_\alpha(H_2)$  for  $\alpha > 0$ .*

*Proof.* Let  $d_1 = d_{H_1}(v)$  and  $d_u = d_{H_1}(u)$  for vertex of  $H_1$  or  $H_2$  that is also a vertex of  $H$ . Then  $d_1 \geq 3$ . Now, we consider the following three cases.

(i) Let  $a_1 = a_2 = 1$ . Then

$$\begin{aligned} SO_\alpha(H_1) - SO_\alpha(H_2) &= 2(d_1^2 + 1)^{\alpha/2} + \sum_{uv \in E(H)} (d_1^2 + d_u^2)^{\alpha/2} - 5^{\alpha/2} \\ &\quad - [(d_1 - 1)^2 + 2^2]^{\alpha/2} - \sum_{uv \in E(H)} [(d_1 - 1)^2 + d_u^2]^{\alpha/2} \\ &= [(d_1^2 + 1)^{\alpha/2} - [d_1^2 + 5 - 2d_1]^{\alpha/2}] + [(d_1^2 + 1)^{\alpha/2} - 5^{\alpha/2}] \\ &\quad + \sum_{uv \in E(H)} [(d_1^2 + d_u^2)^{\alpha/2} - [(d_1 - 1)^2 + d_u^2]^{\alpha/2}] \\ &> 0 \end{aligned}$$

(ii) Let  $a_1 \geq 2$  and  $a_2 = 1$ . Then

$$\begin{aligned} SO_\alpha(H_1) - SO_\alpha(H_2) &= (d_1^2 + 1)^{\alpha/2} - 8^{\alpha/2} + (d_1^2 + 4)^{\alpha/2} - [(d_1 - 1)^2 + 4]^{\alpha/2} \\ &\quad + \sum_{uv \in E(H)} [(d_1^2 + d_u^2)^{\alpha/2} - [(d_1 - 1)^2 + d_u^2]^{\alpha/2}] > 0 \end{aligned}$$

(iii) Let  $a_1 \geq 2$  and  $a_2 \geq 2$ . Then

$$SO_\alpha(H_1) - SO_\alpha(H_2) = 2(d_1^2 + 1)^{\alpha/2} + 5^{\alpha/2} - [(d_1 - 1)^2 + 4]^{\alpha/2} - 2 \times 8^{\alpha/2} + \sum_{uv \in E(H)} \left[ (d_1^2 + d_u^2)^{\alpha/2} - [(d_1 - 1)^2 + d_u^2]^{\alpha/2} \right] > 0$$

Hence  $SO_\alpha(H_1) > SO_\alpha(H_2)$  for  $\alpha > 0$ . □

We recall few basic definitions that are needed for the next section. A graph  $G$  is said to be  $r$ -regular if  $d_G(u) = r$  for all vertices  $u \in V(G)$  and it is said to be *bi-degreed* if it has two distinct vertex degrees. A connected graph  $G$  is said to be *bi-regular* or *semi-regular bipartite* if  $G$  is a bipartite graph with two partite sets  $A$  and  $B$  assuming minimal degree for each vertex in  $A$  and maximal degree for each vertex in  $B$ . Now, we state necessary and sufficient conditions for non-regular graphs to be bi-regular which is proven in [15].

**Lemma 2.** [15] *Let  $G$  be a connected non-regular graph. Then the following statements are equivalent.*

- (i)  $G$  is bi-regular.
- (ii)  $G$  is bi-degreed and the absolute values of  $d_G(u) - d_G(v)$  is a non-zero constant for all edges  $uv$  of  $G$ .
- (iii)  $d_G(u)^2 + d_G(v)^2 > 0$  is constant for all edges  $uv$  of  $G$ .

We present the following inequality which we require in the next section.

**Lemma 3.**  $10^a + 8^a - 13^a - 5^a \geq 0$  for  $0 < a \leq 1$ .

*Proof.* Notice that  $10^a + 8^a - 13^a - 5^a = 0$  if  $a = 1$ . Let  $0 < a < 1$ . Consider the function

$$f(x, y) = 2^a x^a + (x + y)^a - (x + y + 5)^a - x^a, \quad \forall x \geq 1, y \geq 0.$$

Then

$$\frac{\partial f}{\partial x} = a[x^{a-1}(2^a - 1) + (x + y)^{a-1} - (x + y + 5)^{a-1}] > 0$$

and

$$\frac{\partial f}{\partial y} = a[(x + y)^{a-1} - (x + y + 5)^{a-1}] > 0.$$

Thus

$$10^a + 8^a - 13^a - 5^a = f(5, 3) > f(5, 0) = 0.$$

□

### 3. Main Results

In this section, we determine the extremal trees for the general Sombor index  $SO_\alpha$  when  $\alpha > 0$ . We further present the trees with the second minimum general Sombor index  $SO_\alpha$  when  $\alpha > 0$ . First, we prove that the maximum and minimum general Sombor index  $SO_\alpha$  when  $\alpha > 0$  is attained by the star and path respectively.

#### 3.1. Extremal trees for $SO_\alpha$ when $\alpha > 0$ .

**Theorem 1.** *Let  $T$  be a tree of order  $n \geq 4$ . If  $\alpha > 0$ , then*

$$2 \times 5^{\alpha/2} + (n - 3)8^{\alpha/2} \leq SO_\alpha(T) \leq (n - 1)(n^2 - 2n + 2)^{\alpha/2}$$

where the left equality holds if and only if  $T = P_n$ ; and the right equality holds if and only if  $T = S_n$ .

*Proof.* Notice that  $SO_\alpha(P_n) = 2 \times 5^{\alpha/2} + (n - 3)8^{\alpha/2}$ . Now if  $T \neq P_n$ , then by Lemma 1 we can conclude that  $SO_\alpha(T) > SO_\alpha(P_n)$  for  $\alpha > 0$ .

To prove the upper bound, we note that  $d_T(u) + d_T(v) \leq n$  for all edges  $uv$  of  $T$ . Moreover,  $d_T(u)^2 + d_T(v)^2 \leq 1 + (n - 1)^2$  for all edges  $uv$  of  $T$ . Thus


$$SO_\alpha(T) \leq \sum_{uv \in E(T)} [1 + (n - 1)^2]^{\alpha/2} = (n - 1)(n^2 - 2n + 2)^{\alpha/2}$$

where the equality holds if and only if  $d_T(u)^2 + d_T(v)^2 = 1 + (n - 1)^2$  i.e., by Lemma 2,  $T$  is bi-regular. Notice that a tree that is also bi-regular is a star. Hence, the maximal general Sombor index  $SO_\alpha(T)$  is attained by the star  $S_n$  when  $\alpha > 0$ . This completes the proof. □

#### 3.2. Trees with second minimal $SO_\alpha$ when $0 < \alpha \leq 2$ .


Finally, we determine the trees with the second minimum general Sombor index  $SO_\alpha$  when  $0 < \alpha \leq 2$ .

**Remark 1.** For  $n = 5$ , there are only three nonisomorphic trees:  $P_5$ ,  $S_5$  and the tree  $T$  as shown in Fig. 1 (and thus  $T$  has the second minimum general Sombor index).



**Figure 1.** 5-vertex tree  $T$  with the second minimum general Sombor index.


**Remark 2.** For  $n = 6$ , apart from  $P_6$  and  $S_6$ , we have four nonisomorphic trees as shown in Fig. 2.



**Figure 2.** 6-vertex tree with the second minimum general Sombor index.

It can be easily seen with direct computation that the second minimum general Sombor index is attained by the tree  $T_2$  shown in Fig. 2.

Let  $n \geq 7$  and  $s \geq 2$ . Let  $T_{rs}$  denotes the  $n$ -vertex tree with a maximal degree vertex 3 such that  $r + s = n$  and if  $P_r = v_1v_2 \dots v_{r-1}v_r$  is a longest path in  $T_{rs}$  with  $d_{T_{rs}}(v_j) = 3$  for  $3 \leq j \leq r - 2$ , then deleting  $v_j$  we get three paths  $v_1v_2 \dots v_{j-1}$ ,  $v_{j+1} \dots v_{r-1}v_r$  and  $P_s = u_1 \dots u_{s-1}u_s$ . The tree  $T_{rs}$  is shown in Fig. 3 below.



**Figure 3.**  $T_{rs}$

Notice that

$$SO_\alpha(T_{rs}) = 3 \times 5^{\alpha/2} + (n - 7)8^{\alpha/2} + 3 \times 13^{\alpha/2}.$$

We now prove that the second minimum general Sombor index  $SO_\alpha$  is attained by the tree  $T_{rs}$  when  $0 < \alpha \leq 2$ . We first give an outline of the proof.

*Outline of the proof:* We noticed above that the path  $P_n$  has the least general Sombor index. So, we consider any  $n$ -vertex tree  $T$ , which is not  $P_n$  and  $T_{rs}$ , and we proof the theorem by way of contradiction. Such consideration of trees are done based on the degree of the vertex along the longest path in the tree. More precisely, if  $P_r = v_1v_2 \dots v_{r-1}v_r$  is a longest path in  $T$ , then we proceed our argument on the basis of the degree of  $v_2$  or  $v_{r-1}$ .

We assume that the tree  $T$  has the second minimum general Sombor index. Considering (upto isomorphism) different possibilities of  $T$ , we could produce another tree (which could also be  $T_{rs}$ ) with lesser general Sombor index contradicting our assumption that  $T$  has second minimum general Sombor index. Such trees are constructed by attaching only the pendant vertices at some vertices. These are sufficient to conclude our argument as one can notice that the trees with a combination of pendant and non-pendant vertices could also be similarly argued. We make this point clear at one place in *Case 1* of the proof below.

**Theorem 2.** *Let  $T$  be a tree of order  $n \geq 7$  and  $0 < \alpha \leq 2$ . If  $T \neq P_n$ , then*


$$SO_\alpha(T) \geq 3 \times 5^{\alpha/2} + (n - 7)8^{\alpha/2} + 3 \times 13^{\alpha/2}$$

where equality holds if and only if  $T = T_{rs}$ .

*Proof.* Let  $T \neq P_n$  and  $T \neq T_{rs}$  be an  $n$ -vertex tree with the second minimum  $SO_\alpha$  when  $0 < \alpha \leq 2$ . Let  $P_r = v_1v_2 \dots v_{r-1}v_r$  be a path in  $T$  of largest length and let  $d(v) = d_T(v)$  for  $v \in V(T)$ . We consider the following three cases depending on the degree of  $v_2$  or  $v_{r-1}$ .

**Case 1.** Let  $d(v_2) = 2 = d(v_{r-1})$ .

Since  $T \neq P_n$  and  $T \neq T_{rs}$  there exists a vertex  $v_j$  for some  $3 \leq j \leq r - 2$  such that  $d(v_j) \geq 4$  as shown in the left of Fig. 4. Notice that  $u_1, \dots, u_s$  are all pendant vertices. We delete edges  $u_2v_j, \dots, u_s v_j$  ( $s \geq 2$ ) from  $T$  and add them as shown on the right of Fig. 4 to get a new tree  $T'$ . Notice that even if some of the  $u_i$ 's are non-pendant, the construction of  $T'$  is similar (i.e., attaching the path  $u_1u_2 \dots u_s$  to  $v_j$ ) and thus the effect on  $SO_\alpha(T) - SO_\alpha(T')$  remains unchanged.



**Figure 4.** The trees  $T$  (left) and  $T'$  (right), respectively.

Then

$$\begin{aligned} SO_\alpha(T) - SO_\alpha(T') &= [d(v_{j-1})^2 + (s + 2)^2]^{\alpha/2} + [d(v_{j+1})^2 + (s + 2)^2]^{\alpha/2} \\ &\quad + s[1 + (s + 2)^2]^{\alpha/2} - [d(v_{j-1})^2 + 9]^{\alpha/2} - [d(v_{j+1})^2 + 9]^{\alpha/2} \\ &\quad - 13^{\alpha/2} - (s - 2)8^{\alpha/2} - 5^{\alpha/2} \\ &= [d(v_{j-1})^2 + (s + 2)^2]^{\alpha/2} - [d(v_{j-1})^2 + 9]^{\alpha/2} \\ &\quad + [d(v_{j+1})^2 + (s + 2)^2]^{\alpha/2} - [d(v_{j+1})^2 + 9]^{\alpha/2} \end{aligned}$$

$$\begin{aligned}
 &+ (s - 2) \left[ [1 + (s + 2)^2]^{\alpha/2} - 8^{\alpha/2} \right] \\
 &+ 2[1 + (s + 2)^2]^{\alpha/2} - 13^{\alpha/2} - 5^{\alpha/2}
 \end{aligned}$$

Since  $s \geq 2$  notice that  $SO_\alpha(T) - SO_\alpha(T') > 0$ , which is a contradiction to our assumption that  $T$  has the second minimum  $SO_\alpha$ .

**Case 2.** Let  $d(v_2) = 3$  or  $d(v_{r-1}) = 3$ .

Without loss of generality, let  $d(v_2) = 3$ . Following the argument as in *Case 1*, it is enough to consider the following two subcases (involving pendant vertices).

**Subcase 2.1.** Since  $T \neq P_n$  and  $T \neq T_{rs}$  we consider the subcase when  $d(v_j) = 2$  for all  $3 \leq j \leq r - 2$ . Let  $T$  be the tree as shown on the left of Fig. 5. We delete the edge  $v_2u$  from  $T$  and add it to the vertex  $v_3$  to get a new tree  $T'$  as shown on the right of Fig. 5.




**Figure 5.** The trees  $T$  (left) and  $T'$  (right), respectively.

Then

$$SO_\alpha(T) - SO_\alpha(T') = 10^{\alpha/2} + 8^{\alpha/2} - 5^{\alpha/2} - 13^{\alpha/2}.$$

Thus by Lemma 3 we have  $SO_\alpha(T) \geq SO_\alpha(T')$  for  $0 < \alpha \leq 2$ , which is a contradiction to our assumption that  $T$  has the second minimum  $SO_\alpha$ .

**Subcase 2.2.** Since  $T \neq P_n$  and  $T \neq T_{rs}$  there exists a vertex  $v_j$  for some  $3 \leq j \leq r - 2$  such that  $d(v_j) \geq 3$ . The tree  $T$  is shown on the left of Fig. 6. We delete the edges  $uv_2, u_2v_j, \dots, u_s v_j$  and add them as shown on the right of Fig. 6 to get a new tree  $T'$ .



**Figure 6.** The trees  $T$  (left) and  $T'$  (right), respectively.

Then

$$\begin{aligned}
 SO_\alpha(T) - SO_\alpha(T') &= 2 \times 10^{\alpha/2} + [9 + d(v_3)^2]^{\alpha/2} + [d(v_{j-1})^2 + (s + 2)^2]^{\alpha/2} \\
 &+ [d(v_{j+1})^2 + (s + 2)^2]^{\alpha/2} + s[1 + (s + 2)^2]^{\alpha/2} - 2 \times 5^{\alpha/2}
 \end{aligned}$$


$$\begin{aligned}
 & - [4 + d(v_3)^2]^{\alpha/2} - [d(v_{j-1})^2 + 9]^{\alpha/2} - [d(v_{j+1})^2 + 9]^{\alpha/2} \\
 & - 13^{\alpha/2} - (s - 1)8^{\alpha/2} \\
 & = [d(v_{j-1})^2 + (s + 2)^2]^{\alpha/2} - [d(v_{j-1})^2 + 9]^{\alpha/2} \\
 & + [d(v_{j+1})^2 + (s + 2)^2]^{\alpha/2} - [d(v_{j+1})^2 + 9]^{\alpha/2} \\
 & + [9 + d(v_3)^2]^{\alpha/2} - [4 + d(v_3)^2]^{\alpha/2} + 2(10^{\alpha/2} - 5^{\alpha/2}) \\
 & + (s - 1) \left[ [1 + (s + 2)^2]^{\alpha/2} - 8^{\alpha/2} \right] + [1 + (s + 2)^2]^{\alpha/2} - 13^{\alpha/2} \\
 & > 0,
 \end{aligned}$$

which is a contradiction to our assumption that  $T$  has the second minimum  $SO_\alpha$ .

**Case 3.** Let  $d(v_2) \geq 4$  or  $d(v_{r-1}) \geq 4$ .

If  $d(v_3) = 1$ , then  $T$  is a star graph in which case it has the maximum general Sombor index. Thus  $SO_\alpha(T) \geq SO_\alpha(T_{rs})$ , which is a contradiction to the choice of  $T$ . Let  $d(v_3) \geq 2$ . Then there exists a vertex  $v_j$  for some  $4 \leq j \leq r - 2$  such that  $d(v_j) \geq 2$ . In this case, it is enough to consider the following two subcases.

**Subcase 3.1.** Let  $d(v_j) = 2$  for all  $3 \leq j \leq r - 2$ . The tree  $T$  is shown on the left of Fig. 7 and we delete the edges  $v_2u_1, \dots, v_2u_s$  and add them at  $v_j$  as shown on the right of Fig. 7 to get a new tree  $T'$ .




**Figure 7.** The trees  $T$  (left) and  $T'$  (right), respectively.

Then

$$\begin{aligned}
 SO_\alpha(T) - SO_\alpha(T') & = (s + 1)[1 + (s + 2)^2]^{\alpha/2} + [4 + (s + 2)^2]^{\alpha/2} + 2 \times 8^{\alpha/2} \\
 & - 2 \times 5^{\alpha/2} - 3 \times 13^{\alpha/2} - (s - 1)8^{\alpha/2} \\
 & > (s - 3)[(1 + (s + 2)^2)^{\alpha/2} - 8^{\alpha/2}] + 2 \times [(1 + (s + 2)^2)^{\alpha/2} - 5^{\alpha/2}] \\
 & + 3 \times [(1 + (s + 2)^2)^{\alpha/2} - 13^{\alpha/2}] \\
 & > 0,
 \end{aligned}$$

which is a contradiction to our assumption that  $T$  has the second minimum  $SO_\alpha$ .

*Subcase 3.2.* Let  $d(v_j) \geq 3$ . Then the tree  $T$  is as shown on the left of Fig. 8 and we delete the edges  $v_2u_1, \dots, v_2u_s$  and add them at  $v_j$  as shown on the right of Fig. 8 to get a new tree  $T'$ . A simple computation get us that  $SO_\alpha(T) \geq SO_\alpha(T')$ , which is a contradiction to the choice of  $T$ .



**Figure 8.** The trees  $T$  (left) and  $T'$  (right), respectively.

Thus, considering all the cases above, we can conclude that the second minimum general Sombor index tree is attained by the tree  $T_{r_s}$  and this completes the proof.  $\square$

### 4. Conclusion

In this paper, we obtain bounds for the general Sombor index of trees. We further determine the trees with the extremal general Sombor index  $SO_\alpha$  when  $\alpha > 0$ . More precisely, for  $\alpha > 0$  we prove that the path and the star graph have the minimum and the maximum general Sombor index respectively. Further, we determine the tree with the second minimum general Sombor index  $SO_\alpha$  when  $0 < \alpha \leq 2$ .

Following the same argument as in Theorem 1, we note that the lower bound for the general Sombor index is attained by the star when  $\alpha < 0$ . Notice that

$$(n - 1)(n^2 - 2n + 2)^{\alpha/2} \leq SO_\alpha(T)$$

for a tree  $T$  of order  $n \geq 4$  and  $\alpha < 0$ , where the left equality holds if and only if  $T = S_n$ . Determining the extremal trees for the general Sombor index for  $\alpha < 0$  is an attractive future work.

**Acknowledgements.** The authors would like to express their sincere gratitude to the referee(s) for going through the initial manuscript so carefully and for the suggestions made which helped the presentation of the paper.

**Conflict of Interest:** The authors declare that they have no conflict of interest.

**Data Availability:** Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

### References

[1] B. Bollobás and P. Erdős, *Graphs of extremal weights*, *Ars Combin.* **50** (1998), 225–233.

- [2] R. Cruz, I. Gutman, and J. Rada, *Sombor index of chemical graphs*, Appl. Math. Comput. **399** (2021), Article ID: 126018.  
<https://doi.org/10.1016/j.amc.2021.126018>.
- [3] Q. Cui and L. Zhong, *The general Randić index of trees with given number of pendent vertices*, Appl. Math. Comput. **302** (2017), 111–121.  
<https://doi.org/10.1016/j.amc.2017.01.021>.
- [4] ———, *On the general sum-connectivity index of trees with given number of pendent vertices*, Discrete Appl. Math. **222** (2017), 213–221.  
<https://doi.org/10.1016/j.dam.2017.01.016>.
- [5] K.C. Das, A.S. Çevik, I.N. Cangul, and Y. Shang, *On sombor index*, Symmetry **13** (2021), no. 1, Article ID: 140.  
<https://doi.org/10.3390/sym13010140>.
- [6] Z. Du, B. Zhou, and N. Trinajstić, *On the general sum-connectivity index of trees*, Appl. Math. Lett. **24** (2011), no. 3, 402–405.  
<https://doi.org/10.1016/j.aml.2010.10.038>.
- [7] I. Gutman, *Geometric approach to degree-based topological indices: Sombor indices*, MATCH Commun. Math. Comput. Chem. **86** (2021), no. 1, 11–16.
- [8] Y. Hu, X. Li, and Y. Yuan, *Trees with maximum general Randić index*, MATCH Commun. Math. Comput. Chem. **52** (2004), 129–146.
- [9] ———, *Trees with minimum general Randić index*, MATCH Commun. Math. Comput. Chem. **52** (2004), 119–128.
- [10] X. Li, J. Liu, and L. Zhong, *Trees with a given order and matching number that have maximum general Randić index*, Discrete Math. **310** (2010), no. 17-18, 2249–2257.  
<https://doi.org/10.1016/j.disc.2010.04.028>.
- [11] H. Liu, I. Gutman, L. You, and Y. Huang, *Sombor index: review of extremal results and bounds*, J. Math. Chem. **60** (2022), no. 5, 771–798.  
<https://doi.org/10.1007/s10910-022-01333-y>.
- [12] H. Liu, X. Yan, and Z. Yan, *Bounds on the general randić index of trees with a given maximum degree*, MATCH Commun. Math. Comput. Chem. **58** (2007), no. 1, 155–166.
- [13] S. Nikolić, G. Kovačević, A. Miličević, and N. Trinajstić, *The Zagreb indices 30 years after*, Croat. Chem. Acta **76** (2003), no. 2, 113–124.
- [14] C. Phanjoubam and S.M. Mawiong, *On Sombor index and some topological indices*, Iranian J. Math. Chem. **12** (2021), no. 4, 209–215.  
<https://doi.org/10.22052/ijmc.2021.243137.1588>.
- [15] C. Phanjoubam, S.M. Mawiong, and A.M. Buhphang, *On general Sombor index of graphs*, Asian-Eur. J. Math. **16** (2023), no. 3, Article ID: 2350052.  
<https://doi.org/10.1142/S1793557123500523>.
- [16] M. Randić, *Characterization of molecular branching*, J. Am. Chem. Soc. **97** (1975), no. 23, 6609–6615.  
<https://doi.org/10.1021/ja00856a001>.
- [17] I. Redžepović, *Chemical applicability of Sombor indices*, J. Serb. Chem. Soc. **86** (2021), no. 5, 445–457.

- <http://dx.doi.org/10.2298/JSC201215006R>.
- [18] I. Tomescu and M.K. Jamil, *Maximum general sum-connectivity index for trees with given independence number*, MATCH Commun. Math. Comput. Chem. **72** (2014), no. 3, 715–722.
- [19] Z. Wang, Y. Mao, Y. Li, and B. Furtula, *On relations between sombor and other degree-based indices*, J. Appl. Math. Comput. **68** (2022), no. 1, 1–17.  
<https://doi.org/10.1007/s12190-021-01516-x>.
- [20] H. Wiener, *Structural determination of paraffin boiling points*, J. Am. Chem. Soc. **69** (1947), no. 1, 17–20.  
<https://doi.org/10.1021/ja01193a005>.
- [21] L.Z. Zhang, M. Lu, and F. Tian, *Maximum Randić index on trees with  $k$ -pendant vertices*, J. Math. Chem. **41** (2007), no. 2, 161–171.  
<https://doi.org/10.1007/s10910-006-9066-0>.
- [22] L. Zhong, *General Randić index on trees with a given order diameter*, MATCH Commun. Math. Comput. Chem. **62** (2009), no. 1, 177–187.
- [23] L. Zhong and Q. Qian, *The minimum general sum-connectivity index of trees with given matching number*, Bull. Malays. Math. Sci. Soc. **43** (2020), no. 2, 1527–1544.  
<https://doi.org/10.1007/s40840-019-00755-3>.
- [24] B. Zhou and N. Trinajstić, *On a novel connectivity index*, J. Math. Chem. **46** (2009), no. 4, 1252–1270.  
<https://doi.org/10.1007/s10910-008-9515-z>.
- [25] ———, *On general sum-connectivity index*, J. Math. Chem. **47** (2010), no. 1, 210–218.  
<https://doi.org/10.1007/s10910-009-9542-4>.