

Characterizing arc-colored digraphs with an Eulerian trail with restrictions in the color transitions

Carlos Vilchis-Alfaro^{*}, Hortensia Galeana-Sánchez[†]

Instituto de Matemáticas, Universidad Nacional Autónoma de México, CDMX, México

^{*}vilchiscarlos@ciencias.unam.mx

[†]hgaleana@matem.unam.mx

Received: 14 June 2024; Accepted: 30 January 2025

Published Online: 3 February 2025

Abstract: Let H be a digraph possibly with loops, and D a multidigraph without loops. An H -coloring of D is a function $c : A(D) \rightarrow V(H)$. We say that D is an H -colored multidigraph whenever we are taking a fixed H -coloring of D . A trail $W = (v_0, e_0, v_1, e_1, v_2, \dots, v_{n-1}, e_{n-1}, v_n)$ in D is an H -trail if and only if $(c(e_i), c(e_{i+1}))$ is an arc in H , for each $i \in \{0, \dots, n-2\}$. We say that an H -colored multidigraph is H -trail-connected if and only if there is an H -trail starting with arc f_1 and ending with arc f_2 , for any pair of arcs f_1 and f_2 in D . Let D be an H -colored multidigraph and u a vertex of D , the auxiliary digraph D_u is the digraph of allowed transition throughout u .

In this paper we give the following characterization: Let D be an H -colored multidigraph such that the underlying graph of D_u is a disjoint union of complete bipartite graphs, for every $u \in V(D)$. Then D has a Euler H -trail if and only if D is H -trail-connected and, for every $u \in V(D)$, the underlying graph of D_u has a perfect matching. As a consequence we obtain the well-known characterization of the 2-arc-colored multidigraphs containing properly colored Euler trail. Finally, we give an infinite family of digraphs H such that for every multidigraph D without isolated vertices, and every H -coloring of D , the underlying graph of D_u is a disjoint union of complete bipartite graphs and, possibly, isolated vertices, for every $u \in V(D)$.

Keywords: Euler trails, arc-colored digraphs, forbidden transitions

AMS Subject Classification: 05C45, 05C15

1. Introduction

For basic concepts, terminology and notation not defined here, we refer the reader to [3] and [4]. Throughout this work, we will consider finite directed graphs, and

^{*} *Corresponding Author*

directed multigraphs (directed graphs allowing parallel arcs). Let D be a directed graph, $V(D)$ and $A(D)$ will denote the sets of vertices and arcs of D , respectively.

A *(directed) spanning circuit* in a (directed) graph G is defined as a closed (directed) trail that contains each vertex of G . We will say that a (directed) graph is *supereulerian* if it contains a (directed) spanning circuit. Let T be a (directed) spanning circuit in G : if T visits each vertex of G exactly once, then T will be called *Hamilton cycle*; and if T visits each edge of G , then T will be called *Euler trail*. We will say that a (directed) graph is *hamiltonian* if it has a Hamilton cycle and *eulerian* if it contains a Euler trail.

The following graphs were introduced by Harary and Nash-Williams in [12]. Let G be a graph with p vertices and q edges, for $n \geq 2$, $L_n(G)$ will denote the graph with nq vertices that are obtained as follows: for each edge $e = uv$ of G , we take two vertices $f(e, u)$ and $f(e, v)$ in $L_n(G)$ and adding a path with $n - 2$ new intermediate vertices connecting $f(e, u)$ and $f(e, v)$; finally, for each vertex u of G , we add an edge joining $f(e, u)$ and $f(g, u)$, whenever e and g are distinct edges with end point u . They also proved the following relationships between eulerian graphs and hamiltonian cycles of $L_n(G)$.

Theorem 1 ([12]). *Let G be a graph. The following assertions hold:*

1. *If G is eulerian, then $L_n(G)$ is hamiltonian, for every $n \geq 2$.*
2. *If $L_n(G)$ is hamiltonian, for some $n \geq 3$, then G is eulerian.*
3. *G is superulerian if and only if $L_2(G)$ is hamiltonian.*

For a digraph D , the *line digraph* of D , denoted by $L(D)$, is the digraph with vertex set $A(D)$ and $A(L(D)) = \{(a, b) \mid a, b \in V(L(D)), \text{ the head of } a \text{ coincides with the tail of } b\}$. Kasteleyn in [14] show that there is a one-to-one correspondence between the closed directed trails of D and the directed cycles of $L(D)$.

Theorem 2 ([14]). *There is a one-to-one correspondence between the set of closed directed trails in D and the set of directed cycles in $L(D)$.*

Imori et al. [13] introduced the notion of pancircularity as follows. A digraph D with q arcs is said to be *pancircular*, if D contains closed trails of every length L for all $2 \leq L \leq q$.

Corollary 1 ([13]). *Let D be a digraph. Then, D is pancircular if and only if $L(D)$ is pancyclic.*

A graph G is called *edge-colored* if each edge has an assigned color. A *properly colored (PC) walk* is a walk in which no two consecutive edges have the same color, including the last and first edges in a closed walk. Several authors have worked with this

concept, for example, Bang-Jensen et al. [2]. Properly colored walks are of interest as a generalization of walk in undirected and directed graphs, see [3], as well as, in graph theory applications, for example, in genetic and molecular biology [7, 8, 17, 18], channel assignment in wireless networks [1, 19].

Kotzig [15] gave the following characterization of edge-colored multigraphs containing properly colored closed Euler trail.

Theorem 3 (Kotzig [15]). *Let G be an edge-colored eulerian multigraph. Then G has a properly colored closed Euler trail if and only if $\delta_i(x) \leq \sum_{j \neq i} \delta_j(x)$, where $\delta_i(x)$ is the number of edges with color i incident with x , for each vertex x of G .*

Properly colored directed walk in arc-colored directed graphs have also been studied by many authors, for example, Gutin et al. [11], Carraher and Hartke [5, 6] and Gourvès et al. [10]. In [21, 22], arc-colored directed graphs are used to model conflict resolution.

An arc-colored directed graphs D is *PC trail-connected*, if there is a PC trail starting with arc f_1 and ending with arc f_2 , for any pair of arcs f_1, f_2 in D . Sheng et al. [20] characterized 2-arc-colored directed graphs containing PC Euler trails.

Theorem 4 ([20]). *Let D be a 2-arc-colored directed multigraph. Then D has PC Euler trail if and only if D is PC trail-connected and for every $v \in V(D)$, $d_i^+(v) = d_{3-i}^-(v)$, for $i \in \{1, 2\}$.*

Different kinds of edge-coloring in directed and undirected graphs have been studied, for example, in [16] the arcs of a tournament were colored with the vertices of a poset. We will consider the following edge-coloring. Let H be a directed graph possibly with loops and D a directed graph without loops. An H -coloring of D is a function $c : A(D) \rightarrow V(H)$. We will say that D is an H -colored directed graph, whenever we are taking a fixed H -coloring of D . A directed walk $W = (v_0, e_0, v_1, e_1, \dots, e_{k-1}, v_k)$ in D , where $e_i = (v_i, v_{i+1})$ for every i in $\{0, \dots, k-1\}$, is a *directed H -walk* if $(c(e_0), a_0, c(e_1), \dots, c(e_{k-2}), a_{k-2}, c(e_{k-1}))$ is a directed walk in H , with $a_i = (c(e_i), c(e_{i+1}))$ for every $i \in \{0, \dots, k-2\}$. Let W be a directed H -walk, if W is a directed trail then W will be called *H -trail*.

The concepts of H -coloring and H -walks were introduced, for the first time by Linek and Sands in [16], and have been worked mainly in the context of kernel theory and related topics.

A theoretical reason to study H -walks is that they generalize monochromatic walks and properly colored walks. To see that H -walks generalize properly colored walks, notice that if H is a complete graph without loops, then every H -walk is a properly colored walk. And, if H is a graph that only contains loops, then every H -walk is a monochromatic walk. Also, notice that if $W = (x_0, x_1, \dots, x_n)$ is an H -walk such that $(c(x_0x_1), c(x_1x_2), \dots, c(x_{n-1}x_n))$ is a path in H , then W is a rainbow walk.

A motivation for the study of H -walks, in H -colored digraphs, is their possible applications. For example, suppose that we are working with a communication network, represented by a digraph D , where each vertex represent a connection point, and an arc from one connection point A to another connection point B means a way to send information directly from A to B . Notice that more than one arc from A to B can exist since there could be different ways to send information. Moreover, due to different issues (namely risk; such as, damage, attack, virus, blockage, among many others) some arc transitions may be prohibited. In order to have a robust network against communications faults, it is desired to have communication routes with allowed arc transition. To represent this situation we can color the arcs of D , where $A(D) = \{f_1, \dots, f_m\}$, with colors $\{c_1, \dots, c_m\}$, such that $c(f_1) = c_1$; and we define the digraph H , that will determine what arc transitions are allowed, as follows: the set of vertices of H is the set of colors $\{c_1, \dots, c_m\}$, and we add an arc in H from c_i to c_j whenever the transition from arc f_i to arc f_j is allowed. So, if it is required to send a message from point A to point B through the communication network in the most convenient way, we need to find an H -walk in D from A to B .

Another application can be found in roads, roads can be easily represented with digraphs placing a vertex at the intersection of two streets, and an arc from a vertex v to a vertex u if and only if there is a street from v to u without passing through any other corner. Due to traffic laws, it is not always possible to turn from one street to another. So, if we want to find a walk from one corner to another that respect the traffic law, it is not enough to find a walks in a non-arc-colored digraph. So, we can color the arcs of the digraph as stated above. In this case, the digraph H indicates which turns are allowed by traffic laws.

Galeana-Sánchez et al. [9] studied the problem of the existence of closed Euler H -trails in H -colored undirected graphs. They defined the graph G_u as follows: Let u be a vertex of G ; G_u is the graph such that $V(G_u) = \{e \in E(G) \mid e \text{ is incident with } u\}$, and two different vertices a and b are joining by only one edge in G_u if and only if $c(a)$ and $c(b)$ are adjacent in H .

They also showed necessary and sufficient conditions for the existence of closed Euler H -trails, as follows.

Theorem 5 ([9]). *Let H be a graph possibly with loops and G be an H -colored multi-graph without loops. Suppose that G is Eulerian and G_u is a complete k_u -partite graph, for every u in $V(G)$ and for some k_u in $\mathbb{N}$. Then G has a closed Euler H -trail if and only if $|C_i^u| \leq \sum_{j \neq i} |C_j^u|$ for every u in $V(G)$, where $\{C_1^u, \dots, C_{k_u}^u\}$ is the partition of $V(G_u)$ into independent sets.*

The rest of the paper is organized as follows. Section 2 is devoted to give some notation and terminology, which will be used through the paper. In Section 3, we will study the problem of the existence of Euler H -trail. In order to obtain our results we define the auxiliary digraph $L_n^H(D)$. Then, we will prove that there exists a bijection between the set of closed Euler H -trails in D and the set of directed cycles in $L_2^H(D)$.

As a consequence, D has a closed Euler H -trail if and only if $L_n^H(D)$ is hamiltonian, for every $n \geq 2$. Finally, in Section 4 we give an infinite family of digraphs H such that for every multidigraph D without isolated vertices, and every H -coloring of D , the underlying graph of D_u is a disjoint union of complete bipartite graph and, possibly, isolated vertices, for every $u \in V(D)$.

2. Notation and Terminology

Let D be a multidigraph. If e is an arc and u and v are vertices such that $e = (u, v)$, then e is *incident* from u and to v , we will say that u and v are the tail and the head of e , respectively. If $u = v$, then the arc e is a *loop*. The *out-neighborhood* (*in-neighborhood*) of a vertex u , denoted by $N^+(u)$ ($N^-(u)$), is defined as the set of all arcs with tail (head) u . A digraph D' is a *subdigraph* of D if $V(D') \subseteq V(D)$ and $A(D') \subseteq A(D)$. Let S be a nonempty subset of $V(D)$; the subdigraph of D whose vertex set is S , and whose arc set is the set of those arcs of D that have both ends in S , is called the *subdigraph of D induced by S* , and is denoted by $D[S]$. The *underlying graph* of D is the graph G obtained from D by replacing every arc (u, v) with the edge uv .

A directed walk in a digraph D is a sequence $(v_0, e_0, v_1, e_1, \dots, e_{k-1}, v_k)$, where $e_i = (v_i, v_{i+1})$ for every i in $\{0, \dots, k-1\}$. We will say that the direct walk $(v_0, e_0, v_1, e_1, \dots, e_{k-1}, v_k)$ is closed if $v_0 = v_k$. If $v_i \neq v_j$ for all i and j with $i \neq j$, it is called a directed path. A directed cycle is a closed directed walk $(v_0, e_0, v_1, e_1, \dots, e_{k-1}, v_k, e_k, v_0)$, with $k \geq 2$, such that $v_i \neq v_j$ for all i and j with $i \neq j$. In a digraph D a directed walk in which no arc is repeated is a directed trail. A closed Euler directed trail in a digraph D , is a closed directed trail which traverses each arc of D exactly once. If $W = (v_0, e_0, v_1, e_1, \dots, e_{k-1}, v_k)$ and $W' = (v_k, e_k, v_{k+1}, e_{k+1}, \dots, e_{t-1}, v_t)$ are two directed walks, the directed walk $(v_k, e_{k-1}, \dots, e_1, v_1, e_0, v_0)$, obtained by reversing W , is denoted by W^{-1} and the directed walk $(v_0, e_0, v_1, e_1, \dots, e_{k-1}, v_k, e_k, v_{k+1}, e_{k+1}, \dots, e_{t-1}, v_t)$, obtaining by concatenating W and W' at v_k , is denoted by $W \cup W'$. If there is no confusion we will omit adjective "directed".

A simple graph G is said to be multipartite, if for some positive integer k , there exists a partition $X_1, \dots, X_k$ of $V(G)$, such that X_i is an independent set in G (that is no two vertices of X_i are adjacent) for every i in $\{1, \dots, k\}$, in this case, also G is called k -partite. It said that G is a complete k -partite graph whenever G is k -partite and for every u in X_i and for every v in X_j , with $i \neq j$, we have that u and v are adjacent, denoted by $K_{n_1, \dots, n_k}$ where $|X_i| = n_i$ for every i in $\{1, \dots, k\}$. In the particular case when $k = 2$, the graph G is said to be bipartite graph. Let $G_1, G_2, \dots, G_n$ be graphs, the *disjoint union*, denoted by $\bigcup_{i=1}^n G_i$, of $G_1, G_2, \dots, G_n$ is the graph with vertex set, the disjoint union of $V(G_1), V(G_2), \dots, V(G_n)$ and whose edge set, is the disjoint union of $E(G_1), E(G_2), \dots, E(G_n)$. A matching in a graph G is a subset M of $E(G)$, such that no two elements of M are incident with the same vertex in G . A matching M saturates a vertex v if some edge of M is incident with v . If every vertex of G is




Figure 1. The auxiliary digraph $L_2^H(D)$

saturated, the matching M is said to be perfect. We will need the following results.

Theorem 6 (Hall's Theorem). *A bipartite graph with bipartition (X, Y) has a matching that saturates every vertex in X if and only if $|N(S)| \geq |S|$, for every $S \subseteq X$.*

Lemma 1. *Let M and M' be two perfect matching of a graph G . If $M \cap M' = \emptyset$, then there exists a partition of the vertices of G into even cycles. Moreover, every cycle alternates edges between M and M' .*

3. H -trails

In what follows, H will be a digraph possibly with loops, and D will be a multidigraph without loops.

Throughout the paper we will use the auxiliary digraphs D_u and $L_n^H(D)$, which are defined below.

Definition 1. Let D be an H -colored multidigraph and u be a vertex of D . D_u is the digraph such that $V(D_u) = \{f(e, u) \mid e \in A(D) \text{ and } e \text{ is incident with } u\}$, and two different vertices $f(e, u)$ and $f(g, u)$ are joining by only one arc from $f(e, u)$ to $f(g, u)$ in D_u if and only if $e = (x, u)$ and $g = (u, y)$, for some x and y in $V(D)$, and $(c(e), c(g)) \in A(H)$.

Observation 1. D_u is a bipartite digraph with bipartition (X, Y) , where $X = \{f(e, u) \in V(D_u) \mid e = (x, u) \text{ for some } x \in V(D)\}$ and $Y = \{f(e, u) \in V(D_u) \mid e = (u, y) \text{ for some } y \in V(D)\}$. Moreover, if $(f(e, u), f(g, u)) \in A(D_u)$, then $f(e, u) \in X$ and $f(g, u) \in Y$.

Definition 2. Let D be an H -colored multidigraph with $|A(D)| = q$. For $n \geq 2$, $L_n^H(D)$ is the digraph with nq vertices, obtained as follows: for each arc $e = (u, v)$ of D , we take two vertices $f(e, u)$ and $f(e, v)$ in $L_n^H(D)$, and adding a directed path from $f(e, u)$ to $f(e, v)$ with $n - 2$ new intermediate vertices. And the rest of the arcs of $L_n^H(D)$ are defined as follows: $(f(e, u), f(g, u)) \in A(L_n^H(D))$ if and only if $e = (x, u)$ and $g = (u, y)$, for some x and y in $V(D)$, and $(c(e), c(g)) \in A(H)$.

Notice that the digraph $L_n^H(D)$ can be constructed as follows: take the disjoint union of D_u , for every $u \in V(D)$, and for every $e = (u, v) \in A(D)$, we add a directed path from $f(e, u)$ to $f(e, v)$ with $n - 2$ new intermediate vertices, see Figure 1.

Observation 7. For every $e = (u, v) \in A(D)$, we have that $d^+(f(e, u)) = d^-(f(e, v)) = 1$ in $L_n^H(D)$, for each $n \geq 2$. Moreover, when $n = 2$, we have that $N^+(f(e, u)) = \{f(e, v)\}$ and $N^-(f(e, v)) = \{f(e, u)\}$.

Theorem 8. Let D be an H -colored multidigraph. Then there is a bijection between the set of closed H -trails in D and the set of directed cycles in $L_2^H(D)$.

Proof. Let D be an H -colored multidigraph, $\mathcal{P}$ the set of closed H -trails in D and $\mathcal{C}$ the set of directed cycles in $L_2^H(D)$. Consider $T : \mathcal{P} \rightarrow \mathcal{C}$ defined by $T(x_0, e_0, x_1, e_1, x_2, \dots, x_{n-1}, e_{n-1}, x_n) = (f(e_0, x_0), f(e_0, x_1), f(e_1, x_1), f(e_1, x_2), \dots, f(e_{n-1}, x_{n-1}), f(e_{n-1}, x_n), f(e_0, x_0))$.

Claim 1. T is well-defined.

First we will prove that $T(P) = C \in \mathcal{C}$.

It follows from the definition of $L_2^H(D)$ that $(f(e_i, x_i), f(e_i, x_{i+1}))$ is an arc of $L_2^H(D)$, for every $i \in \{0, \dots, n-1\}$.

Let e_i and e_{i+1} be consecutive arcs in P (if $i = n$, then $e_{i+1} = e_1$). Notice that e_i and e_{i+1} are incident with x_{i+1} . Since P is a closed H -trail, we have that $(c(e_i), c(e_{i+1}))$ is an arc in H and $(f(e_i, x_{i+1}), f(e_{i+1}, x_{i+1})) \in A(L_2^H(D))$. Hence, C is a closed walk in $L_2^H(D)$. Since P is a closed H -trail, it follows that C does not repeat a vertex and C is a cycle, i.e., $T(P) = C \in \mathcal{C}$.

Next, we will prove that if $P_1 = P_2$, then $T(P_1) = T(P_2)$.

Let $P_1 = (x_0, e_0, x_1, e_1, x_2, \dots, x_{n-1}, e_{n-1}, x_n)$ and $P_2 = (y_0, f_0, y_1, f_1, y_2, \dots, y_{n-1}, f_{n-1}, y_n)$ in $\mathcal{P}$ such that $P_1 = P_2$.

Since $P_1 = P_2$, we have that $A(P_1) = A(P_2)$ and there exists $k \in \mathbb{N}$ such that $e_i = f_{i+k(\text{mod } n)}$ (since the arcs are traversed in the same order but the first arc is not necessarily the same). It follows from the definition of T that $V(T(P_1)) = V(T(P_2))$ and their are ordered the same. Then, we have that $T(P_1) = T(P_2)$. Therefore, T is well-defined.

Claim 2. T is injective.

Let P_1 and P_2 in $\mathcal{P}$ such that $P_1 \neq P_2$.

If $A(P_1) \neq A(P_2)$, then $V(T(P_1)) \neq V(T(P_2))$ and $T(P_1) \neq T(P_2)$. Otherwise, since $P_1 \neq P_2$, there exists $\{e_1, e_2\} \subseteq A(P_1) = A(P_2)$ such that $e_1 = (x_1, y_1)$ is the arc preceding $e_2 = (x_2, y_2)$ at P_1 and e_1 is not the arc preceding e_2 at P_2 . Hence, $(f(e_1, y_1), f(e_2, x_2)) \in A(T(P_1))$ and $(f(e_1, y_1), f(e_2, x_2)) \notin A(T(P_2))$. Therefore, $T(P_1) \neq T(P_2)$ and T is injective.

Claim 3. T is surjective.

Let C be a cycle in $L_2^H(D)$. It follows from Observation 2 and the fact that $(f(e, x), f(e, y))$ is in $A(L_2^H(D))$, for every $e = (x, y) \in A(D)$, that C must be of the form $C = (f(e_0, x_0), f(e_0, y_0), f(e_1, x_1), f(e_1, y_1), \dots, f(e_q, x_q), f(e_q, y_q), f(e_0, x_0))$, where $e_i = (x_i, y_i) \in A(D)$.

Consider the sequence $P = (x_0, e_0, x_1, e_1, x_2, \dots, x_q, e_q, x_0)$.

Claim 3.1. P is a closed H -trail in D .

For each $k \in \{1, \dots, q\}$, $(f(e_k, y_k), f(e_{k+1}, x_{k+1})) \in A(C) \subseteq A(L_2^H(D))$ (the subindices are taken modulo $q + 1$). So, it follows from the definition of $L_2^H(D)$ that $y_{k-1} = x_k$ and $(c(e_{k-1}), c(e_k)) \in A(H)$. Hence, P is a closed H -walk in D .

On the other hand, since C is a cycle, it follows that e_i appears just once in P , for every $i \in \{1, \dots, q\}$, and P is a closed H -trail.

It follows from the definition of T that $T(P) = C$. Therefore, T is surjective.

By Claims 2 and 3, we have that T is a bijection. $\square$

The underlying graph of $L_n^H(D)$ and D_x will be denoted by $UL_n^H(D)$ and UD_x , respectively. It follows from the definition of $UL_2^H(D)$ that $M_J = \{f(e, x)f(e, y) \in E(UL_2^H(D)) \mid e = (x, y) \in A(D)\}$ is a perfect matching, that we will call the *joint matching* of $UL_2^H(D)$.

Theorem 9. *Let D be an H -colored multidigraph. There exists a partition of the arcs of D into closed H -trails if and only if UD_x has a perfect matching, for every $x \in V(D)$.*

Proof. Let D be an H -colored multidigraph and M_J the joint matching of $UL_2^H(D)$. Suppose that there exists a partition of the arcs of D into closed H -trails, say $P = \{P_1, \dots, P_k\}$.

It follows from Theorem 8 that $C = \{C_1, \dots, C_k\}$ is a set of cycles in $L_2^H(D)$, where $T(P_i) = C_i$. Since $P = \{P_1, \dots, P_k\}$ is a partition of the arcs of D , we have that $V(C_i) \cap V(C_j) = \emptyset$ (because $A(P_i) \cap A(P_j) = \emptyset$) and $V(L_2^H(D)) = \bigcup_{i=1}^k V(C_i)$ (because $A(D) = \bigcup_{i=1}^k A(P_i)$). Therefore, C is a partition of the vertices of $L_2^H(D)$ into cycles. Moreover, $C' = \{C'_1, \dots, C'_k\}$ is a partition of the vertices of $UL_2^H(D)$ into cycles, where C'_i is the underlying cycle of C_i , for each $C_i \in C$.

By construction of C'_i , we have that C'_i alternate edges between M_J and $E(UL_2^H(D)) \setminus M_J$. Hence, $M = \bigcup_{i=1}^k E(C'_i) \setminus M_J$ is a perfect matching of $UL_2^H(D) \setminus M_J$. Therefore, for every $x \in V(D)$, $M_x = M \cap E(UD_x)$ is a perfect matching of UD_x .

Conversely, suppose that M is a perfect matching of $UL_2^H(D) \setminus M_J$. Then, by Lemma 1, we have that there exists a partition of $UL_2^H(D)$ into even cycles, say $C' = \{C'_1, \dots, C'_k\}$, such that every cycle alternate edges between M and M_J .

Since C'_i alternate edges between M_J and M , C'_i must be of the form $C'_i = (f(e_1, x_1), f(e_1, x_2), f(e_2, x_2), f(e_2, x_3), \dots, f(e_n, x_n), f(e_n, x_1), f(e_1, x_1))$.

It follows from Observation 2 that if $(f(e_1, x_1), f(e_1, x_2)) \in A(L_2^H(D))$, then C'_i is a directed cycle in $L_2^H(D)$. Otherwise, $(f(e_1, x_2), f(e_1, x_1)) \in A(L_2^H(D))$ and $C'_i{}^{-1}$ is a directed cycle in $L_2^H(D)$. And then C_i will be the directed cycle, i.e., $C_i = C'_i$ or $C_i = C'_i{}^{-1}$, for every $i \in \{1, \dots, k\}$.

Let $C = \{C_1, \dots, C_k\}$. Since C' is a partition of the vertices of $UL_2^H(D)$ into cycles, it follows that C is a partition of the vertices of $L_2^H(D)$ into directed cycles.

It follows from Theorem 8 that $P = \{P_1, \dots, P_k\}$ is a set of closed H -trails, where $P_i = T^{-1}(C_i)$. Since C is a partition of the vertices of $L_2^H(D)$, it follows that P is a partition of the arcs of D into closed H -trails. $\square$

Definition 3. Let D be an H -colored multidigraph. The H -line digraph, denoted by $L_1^H(D)$, is the digraph such that $V(L_1^H(D)) = A(D)$, and two different vertices $e = (x, y)$ and $g = (u, v)$ are joining by only one arc from e to g in $L_1^H(D)$ if and only if $y = u$ and $(c(e), c(g)) \in A(H)$.

Notice that $L_1^H(D)$ can be obtained from $L_2^H(D)$ by contracting the arc $(f(e, u), f(e, v))$ (and deleting the corresponding loop), for each $e = (u, v) \in A(D)$, in $L_2^H(D)$.

In view of the Theorem 9, we can conclude the following version of Theorem 1 for H -colored multidigraphs.

Theorem 10. Let D be an H -colored multidigraph. Then,

- a. If $L_n^H(D)$ is hamiltonian, for some $n \geq 1$, then D has a closed Euler H -trail.
- b. D has a closed Euler H -trail if and only if $L_n^H(D)$ is hamiltonian, for every $n \geq 1$.

Definition 4. Let D be an H -colored multidigraph with q arcs. We will say that D is an H -pancircular multidigraph if and only if it contains a closed H -trail of length L , for every $2 \leq L \leq q$.

Corollary 2. Let D be an H -colored multidigraph with q arcs. Then, D is an H -pancircular multidigraph if and only if $L_1^H(D)$ is pancyclic.

An H -colored multidigraph D is H -trail-connected, if and only if there is an H -trail starting with arc f_1 and ending with arc f_2 , for any pair of arcs f_1 and f_2 in D .

Theorem 11. Let D be an H -colored multidigraph such that $UD_u = \bigcup_{i=1}^{k_u} K_{n_i^u, m_i^u}$, for every u in $V(D)$ and some $k_u \geq 1$. Then D has a closed Euler H -trail if and only if D is H -trail-connected and, for every $u \in V(D)$, $n_i^u = m_i^u$ for each $i \in \{1, \dots, k_u\}$.

Proof. Let D be an H -colored multidigraph and $D_u = \bigcup_{i=1}^{k_u} K_{n_i^u, m_i^u}$, for every u in $V(D)$ and some $k_u \geq 1$.

Suppose that D has a closed Euler H -trail. Then, D is H -trail-connected and by Theorem 9, UD_u has a perfect matching, for every $u \in V(D)$, say M_u . Since $UD_u = \bigcup_{i=1}^{k_u} K_{n_i^u, m_i^u}$ and M_u is a perfect matching, we have that $n_i^u = m_i^u$, for every $i \in \{1, \dots, k_u\}$.

Conversely, suppose that D is H -trail-connected and, for every $u \in V(D)$, $n_i^u = m_i^u$ for every $i \in \{1, \dots, k_u\}$. Hence, $UD_u = \bigcup_{i=1}^{k_u} K_{n_i^u, n_i^u}$.

It follows from Theorem 6 that UD_u has a perfect matching, for every $u \in V(D)$. Then, by Theorem 9, D has a partition of the arcs of D into closed H -trails, say $\mathcal{P} = \{P_1, \dots, P_k\}$.

If $k = 1$, then D has a closed Euler H -trail. Otherwise, since D is H -trail-connected, there is an H -trail in D starting with arc $e_1 \in A(P_1)$ and ending with arc $e_2 \in A(D) \setminus A(P_1)$, say Q_1 .

Let $g_1 = (x_1, x_2) \in A(Q_1)$ be the first arc in $A(Q_1) \setminus A(P_1)$ and $g_0 = (x_0, x_1)$ the arc prior to g_1 in Q_1 . Hence, $g_0 \in A(P_1)$ and $(c(g_0), c(g_1)) \in A(H)$. Suppose, without loss of generality, that $g_1 \in A(P_2)$.

Let $P_1 = (x_1, f_1, \dots, x_0, g_0, x_1)$ and $P_2 = (x_1, g_1, x_2, \dots, f_0, x_1)$. By the definition of closed H -trail, we have that $\{(c(g_0), c(f_1)), (c(f_0), c(g_1))\} \subseteq A(H)$. By the definition of D_{x_1} , we have that the arcs $(f(g_0, x_1), f(f_1, x_1))$, $(f(f_0, x_1), f(g_1, x_1))$ and $(f(g_0, x_1), f(g_1, x_1))$ are in $A(D_{x_1})$. So, it follows from the Observation 1 and the fact that $UD_{x_1} = \bigcup_{i=1}^{k_{x_1}} K_{n_i^{x_1}, n_i^{x_1}}$ that $(f(f_0, x_1), f(f_1, x_1)) \in A(D_{x_1})$. Therefore, $(c(f_0), c(f_1)) \in A(H)$ and $T_1 = P_1 \cup P_2 = (x_1, f_1, \dots, x_0, g_0, x_1, g_1, x_2, \dots, f_0, x_1)$ is a closed H -trail.

If $A(T_1) = A(D)$, then T_1 is a closed Euler H -trail. Otherwise, there is an H -trail in D starting with arc $e_3 \in A(T_1)$ and ending with arc $e_4 \in A(D) \setminus A(T_1)$, say Q_2 .

Let $g_3 = (v_1, v_2) \in A(Q_2)$ be the first arc in $A(Q_2) \setminus A(T_1)$ and $g_2 = (v_0, v_1)$ the arc prior to g_3 in Q_2 . Hence, $g_2 \in A(T_1)$ and $(c(g_2), c(g_3)) \in A(H)$. Suppose, without loss of generality, that $g_3 \in A(P_3)$.

Let $T_1 = (v_1, f_3, \dots, g_2, v_1)$ and $P_2 = (v_1, g_3, v_2, \dots, f_2, v_1)$. By the definition of closed H -trail, it follows that $\{(c(g_2), c(f_3)), (c(f_2), c(g_3))\} \subseteq A(H)$. By the definition of D_{v_1} , we have that the arcs $(f(g_2, v_1), f(f_3, v_1))$, $(f(f_2, v_1), f(g_3, v_1))$ and $(f(g_2, v_1), f(g_3, v_1))$ are in $A(D_{v_1})$. It follows from the Observation 1 and the fact that $UD_{v_1} = \bigcup_{i=1}^{k_{v_1}} K_{n_i^{v_1}, n_i^{v_1}}$ that the arc $(f(f_2, v_1), f(f_3, v_1))$ is in $A(D_{v_1})$. Therefore, $(c(f_2), c(f_3)) \in A(H)$ and $T_2 = T_1 \cup P_3 = (v_1, f_3, \dots, g_2, v_1, g_3, v_2, \dots, f_2, v_1)$ is a closed H -trail.

If $A(T_2) = A(D)$, then T_2 is a closed Euler H -trail. Otherwise, we can repeat this procedure and after a finite number of steps we obtain that D has a closed Euler H -trail. $\square$

Let D be an c -arc-colored multidigraph and $v \in V(D)$. We will say that $F_v = \{e \in A(D) \mid e \text{ is incident with } v\}$, and $c(F_v) = \{i \mid \text{there exists } e \in F_v \text{ such that } c(e) = i\}$.

Corollary 3. *Let D be a c -arc-colored multidigraph such that $c(F_v) = \{c_1^v, c_2^v\}$, for*

every $v \in V(D)$. Then, D is PC Euler if and only if D is PC trail-connected and for every $v \in V(D)$, $d_{c_i^v}^+(v) = d_{c_{3-i}^v}^-(v)$, for $i \in \{1, 2\}$.

Proof. Suppose that D is a c -arc-colored multidigraph. Then, D is an H -colored multidigraph, where H is the complete digraph without loops and $V(H) = \{1, 2, \dots, c\}$. Notice that if P is a (closed) trail in D , then P is a PC (closed) trail if and only if P is a (closed) H -trail.

Suppose that $c(F_v) = \{c_1^v, c_2^v\}$, for every $v \in V(D)$. Let u be a vertex in $V(D)$.

Claim 1. $UD_u = K_{n_1, m_1} \cup K_{n_2, m_2}$, where $n_i = d_{c_i^u}^-(u)$ and $m_i = d_{c_{3-i}^u}^+(u)$.

Let $E_u^+ = \{f = (u, x) \in A(D) \mid \text{for some } x \in V(D)\}$ and $E_u^- = \{f = (x, u) \in A(D) \mid \text{for some } x \in V(D)\}$. Hence, $F_{c_i^u}^+ = \{f \in E_u^+ \mid c(f) = c_i^u\}$ and $F_{c_i^u}^- = \{f \in E_u^- \mid c(f) = c_i^u\}$ are independent sets in UD_u , for every $i \in \{1, 2\}$. Moreover, for every $i \in \{1, 2\}$, $fg \in E(UD_u)$ if and only if $f \in F_{c_i^u}^-$ and $g \in F_{c_{3-i}^u}^+$.

Notice that $|F_{c_i^u}^+| = d_{c_i^u}^+(u)$ and $|F_{c_i^u}^-| = d_{c_i^u}^-(u)$. Therefore, $UD_u = K_{n_1, m_1} \cup K_{n_2, m_2}$, where $n_i = d_{c_i^u}^-(u)$ and $m_i = d_{c_{3-i}^u}^+(u)$.

Suppose that D is PC Euler. Then, D is PC trail-connected and D has a closed Euler H -trail. It follows from Theorem 11 and Claim 1 that $d_{c_i^u}^-(u) = d_{c_{3-i}^u}^+(u)$, for every $i \in \{1, 2\}$.

Conversely, supposed that D is PC trail-connected and for every $v \in V(D)$, $d_{c_i^v}^+(v) = d_{c_{3-i}^v}^-(v)$, for $i \in \{1, 2\}$. Since D is PC trail-connected, we have that D is H -trail-connected.

It follows from Claim 1 and $d_{c_i^u}^+(u) = d_{c_{3-i}^u}^-(u)$, for $i \in \{1, 2\}$, that $UD_u = K_{n_1, n_1} \cup K_{n_2, n_2}$. Hence, by Theorem 11, D has a closed Euler H -trail, i.e., D is PC Euler. $\square$

Corollary 4 ([20]). *Let D be a 2-arc-colored multidigraph. Then D is PC Euler if and only if D is PC trail-connected and for every $v \in V(D)$, $d_i^+(v) = d_{3-i}^-(v)$, for $i \in \{1, 2\}$.*

Theorem 12. *Let D be an H -colored multidigraph. Then D is H -trail-connected if and only if $L_2^H(D)$ is strongly connected.*

Proof. Let D be an H -colored multidigraph.

Suppose that D is H -trail-connected. Let $f(e, u)$ and $f(g, x)$ in $V(L_2^H(D))$.

Since D is an H -trail-connected graph, there exists an H -trail such that $e = (u_1, u_2)$ is the first arc and $g = (x_1, x_2)$ is the last arc, say $P = (u_1, e, u_2, e_1, \dots, x_1, g, x_2)$. Notice that $T'(P) = (f(e, u_1), f(e, u_2), f(e_1, u_2), \dots, f(g, x_1), f(g, x_2))$ is a path.

Since $f(e, u)$ and $f(g, x)$ are in $V(L_2^H(D))$, it follows that $(u = u_1 \text{ or } u = u_2)$ and $(x = x_1 \text{ or } x = x_2)$. Therefore, there exists a path from $f(e, u)$ to $f(g, x)$ and $L_2^H(D)$ is strongly connected.

Conversely, suppose that $L_2^H(D)$ is strongly connected. Let $e = (x, y)$ and $g = (u, v)$ in $A(D)$.

Since $L_2^H(D)$ is strongly connected, there exists a path from $f(e, x)$ to $f(g, v)$, say P . It follows from Observation 1 that P must be of the form $P = (f(e, x), f(e, y), f(e_1, y), \dots, f(g, u), f(g, v))$. So, $T'^{-1}(P) = (x, e, y, e_1, \dots, u, g, v)$

is an H -trail such that its first arc is e and its last arc is g . Therefore, D is H -trail-connected. $\square$

The next assertion follows directly from the previous results.

Corollary 5. *Given a 2-arc-colored multidigraph, we can check in polynomial time whether D has a close PC Euler trail.*

4. H -coloring

After reading Theorem 11 arises naturally the following question: when does a multidigraph satisfy the hypothesis that $UD_u = \bigcup_{i=1}^{k_u} K_{n_i^u, m_i^u}$, for every u in $V(D)$? In this section we give an infinite family of digraphs H such that for every multidigraph D without isolated vertices, and every H -coloring of D , $UD_u = \bigcup_{i=1}^{k_u} K_{n_i^u, m_i^u}$, for every $u \in V(D)$.

Recall that $V(D_u)$ is the set of arcs in D incident with u , so in this section we will use a simplified notation for the vertices of D_u , the vertex $f(e, u)$ will denote by e . In the remainder of this section we will denote the empty graph with n vertices by $K_{n,0}$.

Theorem 13. *Let H be a digraph, possibly with loops, such that for every pair of vertices u and v of H , $N_H^+(u) = N_H^+(v)$ or $N_H^+(u) \cap N_H^+(v) = \emptyset$. Then, for every multidigraph D without isolated vertices, and every H -coloring of D , $UD_u = \bigcup_{i=1}^{k_u} K_{n_i^u, m_i^u}$, for every $u \in V(D)$ and for some $k_u \geq 1$.*

Proof. Suppose that H is a digraph, possibly with loops, such that for every pair of vertices u and v of H , $N_H^+(u) = N_H^+(v)$ or $N_H^+(u) \cap N_H^+(v) = \emptyset$.

Let D be an H -colored multidigraph without isolated vertices and $u \in V(D)$.

We define $V_0 = \{e \in V(UD_u) : d_{UD_u}(e) = 0\}$. Notice that $UD_u[V_0] = K_{m,0}$, for a non-negative integer m . Thus, we only need to prove that each component in $UD_u - V_0$ is a complete bipartite graph.

Let B be a component of UD_u with at least two vertices. It follows from Observation 1 that B is a bipartite graph with partite sets $X_B = X \cap V(B)$ and $Y_B = Y \cap V(B)$, where X and Y are the partite sets of D_u defined in Observation 1.

Now we will prove that every vertex in X_B is adjacent to every vertex in Y_B . So, let $x \in X_B$ and $y \in Y_B$. Since B is a component of UD_u , there is an xy -path in B , namely $T = (x = e_1, e_2, \dots, e_{2p} = y)$.

Notice that for every $i \in \{1, \dots, p\}$, $e_{2i-1} \in X_B$ and $e_{2i} \in Y_B$. It follows from the definition of D_u that $(c(e_{2i-1}), c(e_{2i}))$ and $(c(e_{2(i+1)-1}), c(e_{2i}))$ are arcs of H , i.e., $N_H^+(c(e_{2i-1})) \cap N_H^+(c(e_{2(i+1)-1})) \neq \emptyset$. So, by hypothesis, $N_H^+(c(e_{2i-1})) = N_H^+(c(e_{2(i+1)-1}))$. Thus, $N_H^+(c(e_1)) = N_H^+(c(e_{2p-1}))$. Moreover, $c(e_{2p}) \in N_H^+(c(e_{2p-1})) = N_H^+(c(e_1))$. Hence, $x = e_1$ and $y = e_{2p}$ are adjacent in B and B is a complete bipartite graph.

Therefore, every component in $UD_u - V_0$ is a complete bipartite graph and $UD_u = \bigcup_{i=1}^{k_u} K_{n_i^u, m_i^u}$, for every $u \in V(D)$ and for some $k_u \geq 1$. $\square$

Proposition 1. *Let H be a complete digraph with a loop in each vertex. Then, for every multidigraph D without isolated vertices, and every H -coloring of D , UD_u is a complete bipartite graph or an empty graph, for every $u \in V(D)$.*

Proof. Suppose that H is a complete digraph with a loop in each vertex. Let D be an H -colored multidigraph without isolated vertices, and u a vertex of D .

Case 1. $d^-(u) = 0$ or $d^+(u) = 0$.

It follows from Observation 1 that UD_u is an empty graph.

Case 2. $d^-(u) \neq 0$ and $d^+(u) \neq 0$.

Let $X = \{e \in A(D) : e = (x, u) \text{ for some } x \in V(D)\}$ and $Y = \{f \in A(D) : f = (u, x) \text{ for some } x \in V(D)\}$. Since $d^-(u) \neq 0$ and $d^+(u) \neq 0$, we have that $X \neq \emptyset$ and $Y \neq \emptyset$.

Consider $e \in X$ and $f \in Y$. Since H is a complete graph with a loop at each vertex, it follows that $(c(e), c(f))$ is an arc in H . Therefore, (e, f) is an arc in D_u .

It follows by Observation 1 that $V(UD_u) = X \cup Y$. Therefore, UD_u is a complete bipartite graph. $\square$

Corollary 6. *Let H be a digraph such that $d^+(x) = 1$, for every vertex x in $V(H)$. Then, for every multidigraph D without isolated vertices, and every H -coloring of D , $UD_u = \bigcup_{i=1}^{k_u} K_{n_i^u, m_i^u}$, for every $u \in V(D)$ and for some $k_u \geq 1$.*

Proof. Let H be a digraph such that $d^+(x) = 1$, for every vertex x in $V(H)$. Hence, for every pair of vertices, u and v , in H , we have that $N_H^+(u) = N_H^+(v)$ or $N_H^+(u) \cap N_H^+(v) = \emptyset$.

Therefore, by Theorem 13, every multidigraph D without isolated vertices, and every H -coloring of D , $UD_u = \bigcup_{i=1}^{k_u} K_{n_i^u, m_i^u}$, for every $u \in V(D)$ and for some $k_u \geq 1$. $\square$

Corollary 7. *Let H be a digraph such that $d^-(x) = 1$, for every vertex x in $V(H)$. Then, for every multidigraph D without isolated vertices, and every H -coloring of D , $UD_u = \bigcup_{i=1}^{k_u} K_{n_i^u, m_i^u}$, for every $u \in V(D)$ and for some $k_u \geq 1$.*

Corollary 8. *If H is a digraph with only loops, then for every multidigraph D without isolated vertices, and every H -coloring of D , $UD_u = \bigcup_{i=1}^{k_u} K_{n_i^u, m_i^u}$, for every $u \in V(D)$ and for some $k_u \geq 1$.*

Corollary 9. *If H is a cycle, then for every multidigraph D without isolated vertices, and every H -coloring of D , $UD_u = \bigcup_{i=1}^{k_u} K_{n_i^u, m_i^u}$, for every $u \in V(D)$ and for some $k_u \geq 1$.*

Corollary 10. *If H is a path with a loop only in the end vertices, then for every multidigraph D without isolated vertices, and every H -coloring of D , $UD_u = \bigcup_{i=1}^{k_u} K_{n_i^u, m_i^u}$, for every $u \in V(D)$ and for some $k_u \geq 1$.*

Acknowledgements: The authors wish to thank the anonymous referees for many comments which substantially improved the rewriting of this paper.

This work was supported by CONACYT FORDECYT-PRONACES/39570/2020; UNAM DGAPA-PAPIIT IN102320; and CONACYT scholarship for postgraduate studies 782239.

Conflict of Interest: The authors declare that they have no conflict of interest.

Data Availability: Data sharing is not applicable to this article as no data sets were generated or analyzed during the current study.

References

- [1] S.K. Ahuja, *Algorithms for Routing and Channel Assignment in Wireless Infrastructure Networks*, The University of Arizona, 2010.
- [2] J. Bang-Jensen, T. Bellitto, and A. Yeo, *On supereulerian 2-edge-coloured graphs*, *Graphs Combin.* **37** (2021), no. 6, 2601–2620.
<https://doi.org/10.1007/s00373-021-02377-8>.
- [3] J. Bang-Jensen and G.Z. Gutin, *Digraphs: Theory, Algorithms and Applications*, Springer Science & Business Media, 2008.
- [4] J.A. Bondy and U.S.R. Murty, *Graph Theory with Applications*, Macmillan London, 1976.
- [5] J.M. Carraher and S.G. Hartke, *Eulerian circuits with no monochromatic transitions in edge-colored digraphs*, *SIAM J. Discrete Math.* **27** (2013), no. 4, 1924–1939.
<https://doi.org/10.1137/120878732>.
- [6] ———, *Eulerian circuits with no monochromatic transitions in edge-colored digraphs with all vertices of outdegree three*, *SIAM J. Discrete Math.* **31** (2017), no. 1, 190–209.
<https://doi.org/10.1137/140992850>.
- [7] D. Dorninger, *Hamiltonian circuits determining the order of chromosomes*, *Discrete Appl. Math.* **50** (1994), no. 2, 159–168.
[https://doi.org/10.1016/0166-218X\(92\)00171-H](https://doi.org/10.1016/0166-218X(92)00171-H).
- [8] D. Dorninger and W. Timischl, *Geometrical constraints on bennett’s predictions of chromosome order*, *Heredity* **59** (1987), no. 3, 321–325.
<https://doi.org/10.1038/hdy.1987.138>.
- [9] H. Galeana-Sánchez, R. Rojas-Monroy, R. Sánchez-López, and J.I. Villarreal-Valdés, *Some conditions for the existence of euler h -trails*, *Graphs Combin.* **35** (2019), no. 5, 1197–1208.
<https://doi.org/10.1007/s00373-019-02066-7>.
- [10] L. Gourvès, A. Lyra, C.A. Martinhon, and J. Monnot, *Complexity of trails, paths and circuits in arc-colored digraphs*, *Discrete Appl. Math.* **161** (2013), no. 6, 819–

828.
<https://doi.org/10.1016/j.dam.2012.10.025>.
- [11] G. Gutin, B. Sudakov, and A. Yeo, *Note on alternating directed cycles*, Discrete Math. **191** (1998), no. 1-3, 101–107.
[https://doi.org/10.1016/S0012-365X\(98\)00097-1](https://doi.org/10.1016/S0012-365X(98)00097-1).
- [12] F. Harary and C.S.J. Nash-Williams, *On eulerian and hamiltonian graphs and line graphs*, Canad. Math. Bull. **8** (1965), no. 6, 701–709.
<https://doi.org/10.4153/CMB-1965-051-3>.
- [13] M. Imori, M. Matsumoto, and H. Yamada, *The line digraph of a regular and pancircular digraph is also regular and pancircular*, Graphs Combin. **4** (1988), no. 1, 235–239.
<https://doi.org/10.1007/BF01864164>.
- [14] P.W. Kasteleyn, *A soluble self-avoiding walk problem*, Physica **29** (1963), no. 12, 1329–1337.
[https://doi.org/10.1016/S0031-8914\(63\)80241-4](https://doi.org/10.1016/S0031-8914(63)80241-4).
- [15] A. Kotzig, *Moves without forbidden transitions in a graph*, Matematickỳ časopis **18** (1968), no. 1, 76–80.
- [16] V. Linek and B. Sands, *A note on paths in edge-coloured tournaments*, Ars Combin. **44** (1996), 225–228.
- [17] P.A. Pevzner, *Dna physical mapping and alternating eulerian cycles in colored graphs*, Algorithmica **13** (1995), no. 1, 77–105.
<https://doi.org/10.1007/BF01188582>.
- [18] P.A. Pevzner, H. Tang, and M.S. Waterman, *An eulerian path approach to dna fragment assembly*, Proc. Nat. Acad. Sci. **98** (2001), no. 17, 9748–9753.
<https://doi.org/10.1073/pnas.171285098>.
- [19] S. Sankararaman, A. Efrat, S. Ramasubramanian, and P.K. Agarwal, *On channel-discontinuity-constraint routing in wireless networks*, Ad Hoc Netw. **13** (2014), 153–169.
<https://doi.org/10.1016/j.adhoc.2011.04.011>.
- [20] B. Sheng, R. Li, and G. Gutin, *The euler and chinese postman problems on 2-arc-colored digraphs*, arXiv preprint arXiv:1707.06503 (2017).
- [21] H. Xu, D.M. Kilgour, K.W. Hipel, and G. Kemkes, *Using matrices to link conflict evolution and resolution in a graph model*, European J. Oper. Res. **207** (2010), no. 1, 318–329.
<https://doi.org/10.1016/j.ejor.2010.03.025>.
- [22] H. Xu, K.W. Li, D.M. Kilgour, and K.W. Hipel, *A matrix-based approach to searching colored paths in a weighted colored multidigraph*, Appl. Math. Comput. **215** (2009), no. 1, 353–366.
<https://doi.org/10.1016/j.amc.2009.04.086>.