

A Note on Distance-Fall Colorings

Wayne Goddard^{1,*}, Sonwabile Mafunda^{2,3}

¹School of Mathematical and Statistical Sciences Clemson University, USA
*goddard@clemson.edu

²Soka University of America, USA

³University of Johannesburg, South Africa
smafunda@soka.edu

Received: 29 August 2025; Accepted: 22 December 2025
Published Online: 6 January 2025

Dedicated to Odile Favaron

Abstract: We say a proper coloring of a graph is distance- k fall if every vertex is within distance k of at least one vertex of every color. We show that if G is a connected graph of order at least 3 that is 3-colorable, then it has a distance-2 fall 3-coloring. Further, for every integer $k \geq 2$, if T is a tree of order at least k , then T has a k -coloring such that every vertex is within distance $k - 1$ of every color. This proves an old conjecture of Beineke and Henning that every tree of order n has an independent distance- d -dominating set of size at most $n/(d + 1)$.

Keywords: fall coloring, distance- k domination, chromatic number.

AMS Subject classification: 05C12, 05C15, 05C69

1. Introduction

A *fall coloring* is a proper coloring of the vertices of a graph such that every vertex sees every color. That is, for each vertex v its closed neighborhood $N[v]$ contains all colors. Not all graphs have a fall coloring. The simplest example is the 5-cycle. In this paper we consider proper colorings where every vertex is “near” to every color. We say a coloring is *distance- k fall* if every vertex is within distance k of at least one vertex of every color. For example, for a graph of diameter 2, every proper coloring is distance-2 fall; and every odd cycle has a 3-coloring that is distance-2 fall.

* *Corresponding Author*

The terminology “*fall coloring*” was introduced in the 2000 paper by Dunbar et al. [5]. But the concept is older, having been studied before as partitioning a graph into independent dominating sets; see the references of [5]. Recall that a set S of vertices is *independent* if no two elements of S are joined by an edge. A set S is *dominating* if every vertex is either in S or adjacent to at least one vertex of S . An *independent dominating* set is one that is both independent and dominating. More generally, a set S is *distance- k dominating* if every vertex is within distance k of at least one vertex of S . Thus a distance- k fall coloring is a partition of the vertex set into independent distance- k dominating sets. The parameter *independent distance- k domination*, that is, the minimum size of an independent distance- k dominating set, was originally studied by Beineke and Henning [1].

2. Distance-Fall Colorings and Chromatic Number

It is immediate that if a (connected nontrivial) graph is bipartite, then the bipartite coloring is a fall coloring. So the next question to consider is 3-colorable graphs. For a coloring, we say a vertex is *d -good* if every color appears within distance d of the vertex; otherwise the vertex is *d -bad*. The following theorem shows that if the graph is 3-colorable then there is a proper coloring with 3 colors such that every vertex is 2-good.

Theorem 1. *If G is a connected graph of order at least 3 that is 3-colorable, then G has a distance-2 fall 3-coloring.*

Proof. Suppose to the contrary, that G does not admit a distance-2 fall 3-coloring. Take a proper 3-coloring of G that minimizes the number of 2-bad vertices. Without loss of generality, choose a 2-bad vertex $v \in V(G)$. Then $N(v)$ is monochromatic. If there exists a vertex $w_1 \in N(v)$ with $\deg(w_1) = 1$, then by recoloring w_1 with the missing color in $N[v]$, we obtain a proper 3-coloring of G with strictly fewer 2-bad vertices, contradicting the minimality of the original coloring.

Therefore, assume $\deg(w) \geq 2$ for all $w \in N(v)$. Since v is 2-bad, we have $N(v) \cap N(w) = \emptyset$ and $N(w)$ is monochromatic for every $w \in N(v)$. That is, v and all vertices in its second neighborhood share the same color.

Now recolor v with the missing color in $N[v]$. Vertex v then becomes 2-good, and every neighbor of v remains 2-good. Moreover, no previously 2-good vertex becomes 2-bad under this recoloring. Thus, the total number of 2-bad vertices decreases, contradicting the minimality of the original coloring.

Therefore, there exists a proper 3-coloring of G with zero 2-bad vertices. \square

It is perhaps interesting to note that an independent distance-2 dominating set is related to an independent isolating set. An isolating (or vertex-edge dominating) set S can be viewed as a 2-dominating set S with the added condition that if vertex v is at distance 2 from S , then all its neighbors are at distance 1 from S . An independent

isolating set is an isolating set that is also independent. See for example [2, 3, 6]. It is easy to see that Theorem 1 does not extend to a partition into three independent isolating sets (for example C_5). However, in [3] it was shown that it almost does: specifically, if G is 3-colorable there exist three sets A , B , and C of vertices, such that each is an independent isolating set, their union is all vertices, and such that A and B overlap in at most one vertex, while C is disjoint from $A \cup B$.

It is, however, unclear if Theorem 1 generalizes to larger number of colors. The next step would be to determine whether every connected 4-colorable graph of order at least 4 has a 4-coloring such that every vertex is within distance three of all colors. This is best possible because of path-complete graph, meaning the graph that consists of a path and a complete graph joined by an edge. However, in the case of trees, the above theorem does generalize:

Theorem 2. *For integer $k \geq 2$, if T is a tree of order at least k , then T has a k -coloring such that every vertex is within distance $k - 1$ of every color.*

Proof. Let T be a tree of diameter d . If $d < k$, then any coloring that uses each color at least once automatically has the desired property.

So, assume $d \geq k$. Let $P := v_1, v_2, \dots, v_d$ be a diametral path in T . Define a coloring $c : V(P) \rightarrow \{0, 1, 2, \dots, k - 1\}$ on the vertices of P such that $c(v_i) = i \pmod{k}$.

Now, let $e = uv$ be a central edge of P , and let T_u and T_v be the components of $T - e$ containing u and v , respectively. If $T \setminus P$ denotes the forest induced by the vertices of T not contained in P , then color every vertex of $T \setminus P$ such that, in each of the components T_u and T_v , any two vertices at the same distance from u or v receive the same color.

We claim that this coloring has the desired property. If a vertex w lies within distance $(k - 1)/2$ of the edge e , then w is within distance $k - 1$ of each color on P . Otherwise, the unique path from w to P enters P at some vertex and continues along P via e ; the first k vertices on this path all have distinct colors. Hence the theorem follows. \square

Theorem 1 is equivalent to saying that the vertices of a 3-colorable graph can be partitioned into three (disjoint) independent distance-2 dominating sets. And thus the independent distance-2 domination number of a 3-colorable graph is at most $n/3$, where n is the order. This generalizes Theorem 2 of [1], which proved the bound for trees. Furthermore, Theorem 2 shows that the independent distance- k domination number of a tree is at most $n/(k+1)$, which establishes the Conjecture at the end of [1].

We note that Theorem 2 and thus the conjecture in [1] was recently also proved by Bujtás et al. [4]. Indeed, they showed that Theorem 2 generalizes to bipartite graphs.

There is a version of Theorem 1 for general graphs with a slightly weaker distance condition. A *partial coloring* means a proper coloring where only some of the vertices are colored.

Theorem 3. *If G is a connected graph of order at least 3, then G has a partial 3-coloring such that every vertex is within distance 3 of every color.*

Proof. Consider any partial 3-coloring that maximizes the number of colored vertices. Note that any uncolored vertex v has neighbors of every color, since it is not possible to color v . Thus, out of all partial 3-colorings with the maximum number of colored vertices, take the one with the minimum number of 2-bad vertices and let v be such a vertex.

We know v is colored. Furthermore, all of its neighbors are colored. If some vertex w at distance 2 from v is uncolored, then v is 3-good since $N(w)$ contains all colors. So assume every vertex within distance two of v is colored. Then, as in the proof of Theorem 1, one can re-color either v or a neighbor of v so that v and its neighbors are 2-good. Note that if this recoloring makes an uncolored vertex bad, then we have a contradiction of the original requirement that the maximum number of vertices were colored. So every vertex that is 2-bad is 3-good; or in other words, every vertex is 3-good. \square

This strengthens the first result in [1] for the case that $k = 3$.

We conclude with a brief comment about graph operations. Kaul and Mitillos [7] showed that if a graph G has a fall k -coloring and a graph H has a k -coloring, then the cartesian product $G \square H$ has a fall k -coloring. The same idea works here: if G has a distance- d fall k -coloring f_G and H has a k -coloring f_H , then the cartesian product $G \square H$ has a distance- d fall k -coloring. As per usual in the cartesian product, consider both colorings as assigning integers in the range 1 to k and take the coloring of vertex (g, h) in the product to be the sum $f_G(g) + f_H(h)$ modulo k .

We note further that if G has a distance- d_G fall k_G -coloring and H has a distance- d_H fall k_H -coloring, then the cartesian product $G \square H$ has a distance- $(d_G + d_H)$ fall $k_G k_H$ -coloring. Simply take the coloring of vertex (g, h) in the product to be the ordered pair $(f_G(g), f_H(h))$.

Similar results can be shown for other products.

Acknowledgements: The authors would like to thank Ben Gobler for suggesting the concept.

Conflict of Interest: The authors declare that they have no conflict of interest.

Data Availability. Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

References

- [1] L.W. Beineke and M.A. Henning, *Some extremal results on independent distance domination in graphs*, *Ars Combin.* **37** (1994), 223–233.

-
- [2] R. Boutrig, M. Chellali, T.W. Haynes, and S.T. Hedetniemi, *Vertex-edge domination in graphs*, *Aequ. Math.* **90** (2016), no. 2, 355–366.
<https://doi.org/10.1007/s00010-015-0354-2>.
- [3] G. Boyer and W. Goddard, *Bounds on independent isolation in graphs*, *Discrete Appl. Math.* **372** (2025), 143–149.
<https://doi.org/10.1016/j.dam.2025.04.016>.
- [4] C. Bujtás, V.I. Chenoweth, S. Klavžar, and G. Zhang, *Revisiting d -distance (independent) domination in trees and in bipartite graphs*, arXiv preprint arXiv:2508.12804 (2025).
- [5] J.E. Dunbar, S.M. Hedetniemi, S.T. Hedetniemi, D.P. Jacobs, J. Knisely, R.C. Laskar, and D.F. Rall, *Fall colorings of graphs*, *J. Combin. Math. Combin. Comput.* **33** (2000), 257–273.
- [6] O. Favaron and P. Kaemawichanurat, *Inequalities between the K_k -isolation number and the independent K_k -isolation number of a graph*, *Discrete Appl. Math.* **289** (2021), 93–97.
<https://doi.org/10.1016/j.dam.2020.09.011>.
- [7] H. Kaul and C. Mitillos, *On graph fall-coloring—operators and heredity*, *J. Combin. Math. Combin. Comput.* **106** (2018), 125–151.